

Prospects for A Fixed-Target Experiment at the LHC: AFTER@LHC

Jean-Philippe Lansberg
IPN Orsay, Université Paris-Sud

BARYONS 2013, June 24-28 2013



thanks to M. Anselmino (Torino), R. Arnaldi (Torino), S.J. Brodsky (SLAC), V. Chambert (IPNO), J.P. Didelez (IPNO), E.G. Ferreira (USC), F. Fleuret (LLR), B. Genolini (IPNO), C. Hadjidakis (IPNO), C. Lorcé (IPNO), A. Rakotozafindrabe (CEA), P. Rosier (IPNO), I. Schienbein (LPSC), E. Scomparin (Torino), and U.I. Uggerhøj (Aarhus)

Outline

Part 1: Why a new fixed-target experiment for High-Energy Physics now ?

Outline

Part 1: Why a new fixed-target experiment for High-Energy Physics now ?

Part 2: A Fixed-Target Experiment using LHC beams: AFTER@LHC

Outline

Part 1: Why a new fixed-target experiment for High-Energy Physics now ?

Part 2: A Fixed-Target Experiment using LHC beams: AFTER@LHC

Part 3: Flagship studies

Outline

Part 1: Why a new fixed-target experiment for High-Energy Physics now ?

Part 2: A Fixed-Target Experiment using LHC beams: AFTER@LHC

Part 3: Flagship studies

Conclusions and Outlooks

Part I

Why a new fixed-target experiment for High-Energy Physics now ?

Decisive advantages of Fixed-target experiments

- Fixed-target experiments offer specific **advantages** that are still nowadays **difficult to challenge by collider experiments**

Decisive advantages of Fixed-target experiments

- Fixed-target experiments offer specific **advantages** that are still nowadays **difficult to challenge by collider experiments**
- They exhibit 4 decisive features,
 - accessing the **high** Feynman x_F domain ($x_F \equiv \frac{p_z}{p_{z\max}}$)
 - achieving **high luminosities** with dense targets,
 - **varying** the atomic mass of the **target** almost at will,
 - **polarising** the target.

The European strategy for particle physics

Approved by the CERN council at the special Session held in Lisbon on July 14, 2006

The European strategy for particle physics

Approved by the CERN council at the special Session held in Lisbon on July 14, 2006

9. A variety of important research lines are at the interface between particle and nuclear physics requiring dedicated experiments; *Council will seek to work with NuPECC in areas of mutual interest, and maintain the capability to perform **fixed target experiments at CERN.***

pg. 37 of the Strategy Brochure

The European strategy for particle physics

Updated by the CERN council at the special Session held in Brussels on May 30, 2013

- k. A variety of research lines at the boundary between particle and nuclear physics require dedicated experiments. *The CERN Laboratory should maintain its capability to perform **unique experiments**. CERN should continue to work with NuPECC on topics of mutual interest.*

pg 22. of the Strategy Update Brochure

The European strategy for particle physics

Updated by the CERN council at the special Session held in Brussels on May 30, 2013

- k. A variety of research lines at the boundary between particle and nuclear physics require dedicated experiments. *The CERN Laboratory should maintain its capability to perform **unique experiments**. CERN should continue to work with NuPECC on topics of mutual interest.*

pg 22. of the Strategy Update Brochure

Using the LHC beams, for the first time,
the 100-GeV frontier can be broken at a fixed target experiment,

The European strategy for particle physics

Updated by the CERN council at the special Session held in Brussels on May 30, 2013

- k. A variety of research lines at the boundary between particle and nuclear physics require dedicated experiments. *The CERN Laboratory should maintain its capability to perform **unique experiments**. CERN should continue to work with NuPECC on topics of mutual interest.*

pg 22. of the Strategy Update Brochure

Using the LHC beams, for the first time,
the 100-GeV frontier can be broken at a fixed target experiment,

- without affecting the LHC performance
- with an extracted beam line using a bent crystal

The European strategy for particle physics

Updated by the CERN council at the special Session held in Brussels on May 30, 2013

- k. A variety of research lines at the boundary between particle and nuclear physics require dedicated experiments. *The CERN Laboratory should maintain its capability to perform **unique experiments**. CERN should continue to work with NuPECC on topics of mutual interest.*

pg 22. of the Strategy Update Brochure

Using the LHC beams, for the first time,
the 100-GeV frontier can be broken at a fixed target experiment,

- without affecting the LHC performance
- with an extracted beam line using a bent crystal
- with the possibility of polarising the target
- without target-species limitation

The European strategy for particle physics

Updated by the CERN council at the special Session held in Brussels on May 30, 2013

- k. A variety of research lines at the boundary between particle and nuclear physics require dedicated experiments. *The CERN Laboratory should maintain its capability to perform **unique experiments**. CERN should continue to work with NuPECC on topics of mutual interest.*

pg 22. of the Strategy Update Brochure

Using the LHC beams, for the first time,
the 100-GeV frontier can be broken at a fixed target experiment,

- without affecting the LHC performance
- with an extracted beam line using a bent crystal
- with the possibility of polarising the target
- without target-species limitation
- with an outstanding luminosity, yet without pile-up
- with virtually no limit on particle-species studies (except top quark)

The European strategy for particle physics

Updated by the CERN council at the special Session held in Brussels on May 30, 2013

- k. A variety of research lines at the boundary between particle and nuclear physics require dedicated experiments. *The CERN Laboratory should maintain its capability to perform **unique experiments**. CERN should continue to work with NuPECC on topics of mutual interest.*

pg 22. of the Strategy Update Brochure

Using the LHC beams, for the first time,
the 100-GeV frontier can be broken at a fixed target experiment,

- without affecting the LHC performance
- with an extracted beam line using a bent crystal
- with the possibility of polarising the target
- without target-species limitation
- with an outstanding luminosity, yet without pile-up
- with virtually no limit on particle-species studies (except top quark)
- with modern detection techniques

The European strategy for particle physics

Updated by the CERN council at the special Session held in Brussels on May 30, 2013

- k. A variety of research lines at the boundary between particle and nuclear physics require dedicated experiments. *The CERN Laboratory should maintain its capability to perform **unique experiments**. CERN should continue to work with NuPECC on topics of mutual interest.*

pg 22. of the Strategy Update Brochure

Using the LHC beams, for the first time,
the 100-GeV frontier can be broken at a fixed target experiment,

- without affecting the LHC performance
- with an extracted beam line using a bent crystal
- with the possibility of polarising the target
- without target-species limitation
- with an outstanding luminosity, yet without pile-up
- with virtually no limit on particle-species studies (except top quark)
- with modern detection techniques

AFTER@LHC would definitely be a **unique experiment**

Part II

A fixed-target experiment using the LHC beam(s): AFTER@LHC

Generalities

- pp or pA collisions with a 7 TeV p^+ on a fixed target occur at a CM energy

$$\sqrt{s} = \sqrt{2m_N E_p} \simeq 115 \text{ GeV}$$

Generalities

- pp or pA collisions with a 7 TeV p^+ on a fixed target occur at a CM energy

$$\sqrt{s} = \sqrt{2m_N E_p} \simeq 115 \text{ GeV}$$

- In a symmetric collider mode, $\sqrt{s} = 2E_p$, *i.e.* much larger

Generalities

- pp or pA collisions with a 7 TeV p^+ on a fixed target occur at a CM energy

$$\sqrt{s} = \sqrt{2m_N E_p} \simeq 115 \text{ GeV}$$

- In a symmetric collider mode, $\sqrt{s} = 2E_p$, *i.e.* much larger
- Benefit of the fixed target mode : **boost**: $\gamma_{CM}^{Lab} = \frac{\sqrt{s}}{2m_p} \simeq 60$

Generalities

- pp or pA collisions with a 7 TeV p^+ on a fixed target occur at a CM energy

$$\sqrt{s} = \sqrt{2m_N E_p} \simeq 115 \text{ GeV}$$

- In a symmetric collider mode, $\sqrt{s} = 2E_p$, *i.e.* much larger
- Benefit of the fixed target mode : **boost**: $\gamma_{CM}^{Lab} = \frac{\sqrt{s}}{2m_p} \simeq 60$
 - Consider a **photon emitted at 90°** w.r.t. the z-axis (beam) in the CM:
($p_{z,CM} = 0$, $E_{CM}^\gamma = p_T$)

Generalities

- pp or pA collisions with a 7 TeV p^+ on a fixed target occur at a CM energy

$$\sqrt{s} = \sqrt{2m_N E_p} \simeq 115 \text{ GeV}$$

- In a symmetric collider mode, $\sqrt{s} = 2E_p$, *i.e.* much larger
- Benefit of the fixed target mode : **boost**: $\gamma_{CM}^{Lab} = \frac{\sqrt{s}}{2m_p} \simeq 60$
 - Consider a **photon emitted at 90°** w.r.t. the z-axis (beam) in the CM:
 $(p_{z,CM} = 0, E_{CM}^\gamma = p_T)$
 - $$\begin{pmatrix} E_{Lab} \\ p_{z,Lab} \end{pmatrix} = \begin{pmatrix} \gamma & \gamma\beta \\ \gamma\beta & \gamma \end{pmatrix} \begin{pmatrix} p_T \\ 0 \end{pmatrix}$$

Generalities

- pp or pA collisions with a 7 TeV p^+ on a fixed target occur at a CM energy

$$\sqrt{s} = \sqrt{2m_N E_p} \simeq 115 \text{ GeV}$$

- In a symmetric collider mode, $\sqrt{s} = 2E_p$, *i.e.* much larger
- Benefit of the fixed target mode : **boost**: $\gamma_{CM}^{Lab} = \frac{\sqrt{s}}{2m_p} \simeq 60$
 - Consider a **photon emitted at 90°** w.r.t. the z-axis (beam) in the CM: $(p_{z,CM} = 0, E_{CM}^\gamma = p_T)$
 - $\begin{pmatrix} E_{Lab} \\ p_{z,Lab} \end{pmatrix} = \begin{pmatrix} \gamma & \gamma\beta \\ \gamma\beta & \gamma \end{pmatrix} \begin{pmatrix} p_T \\ 0 \end{pmatrix}$
 - $p_{z,Lab} \simeq 60p_T$! [A 67 MeV γ from a π^0 at rest in the CM can easily be detected.]

Generalities

- pp or pA collisions with a **7 TeV p^+** on a fixed target occur at a CM energy

$$\sqrt{s} = \sqrt{2m_N E_p} \simeq 115 \text{ GeV}$$

- In a symmetric collider mode, $\sqrt{s} = 2E_p$, *i.e.* much larger
- Benefit of the fixed target mode : **boost**: $\gamma_{CM}^{Lab} = \frac{\sqrt{s}}{2m_p} \simeq 60$

- Consider a **photon emitted at 90°** w.r.t. the z-axis (beam) in the CM:

$$\begin{pmatrix} E_{Lab} \\ p_{z,Lab} \end{pmatrix} = \begin{pmatrix} \gamma & \gamma\beta \\ \gamma\beta & \gamma \end{pmatrix} \begin{pmatrix} p_T \\ 0 \end{pmatrix} \quad (p_{z,CM} = 0, E_{CM}^\gamma = p_T)$$

- $p_{z,Lab} \simeq 60p_T$! [A 67 MeV γ from a π^0 at rest in the CM can easily be detected.]

- Angle in the Lab. frame: $\tan \theta = \frac{p_T}{p_{z,Lab}} = \frac{1}{\gamma\beta} \Rightarrow \theta \simeq 1^\circ$.

[Rapidity shift: $\Delta y = \tanh^{-1} \beta \simeq 4.8$]

Generalities

- pp or pA collisions with a **7 TeV p^+** on a fixed target occur at a CM energy

$$\sqrt{s} = \sqrt{2m_N E_p} \simeq 115 \text{ GeV}$$

- In a symmetric collider mode, $\sqrt{s} = 2E_p$, *i.e.* much larger
- Benefit of the fixed target mode : **boost**: $\gamma_{CM}^{Lab} = \frac{\sqrt{s}}{2m_p} \simeq 60$
 - Consider a **photon emitted at 90°** w.r.t. the z-axis (beam) in the CM: $(p_{z,CM} = 0, E_{CM}^\gamma = p_T)$
 - $\begin{pmatrix} E_{Lab} \\ p_{z,Lab} \end{pmatrix} = \begin{pmatrix} \gamma & \gamma\beta \\ \gamma\beta & \gamma \end{pmatrix} \begin{pmatrix} p_T \\ 0 \end{pmatrix}$
 - $p_{z,Lab} \simeq 60p_T$! [A 67 MeV γ from a π^0 at rest in the CM can easily be detected.]
- Angle in the Lab. frame: $\tan \theta = \frac{p_T}{p_{z,Lab}} = \frac{1}{\gamma\beta} \Rightarrow \theta \simeq 1^\circ$.
[Rapidity shift: $\Delta y = \tanh^{-1} \beta \simeq 4.8$]
- The entire forward CM hemisphere ($y_{CM} > 0$) within $0^\circ \leq \theta_{Lab} \leq 1^\circ$
[$y_{CM} = 0 \Rightarrow y_{Lab} \simeq 4.8$]

Generalities

- pp or pA collisions with a **7 TeV p^+** on a fixed target occur at a CM energy

$$\sqrt{s} = \sqrt{2m_N E_p} \simeq 115 \text{ GeV}$$

- In a symmetric collider mode, $\sqrt{s} = 2E_p$, *i.e.* much larger
- Benefit of the fixed target mode : **boost**: $\gamma_{CM}^{Lab} = \frac{\sqrt{s}}{2m_p} \simeq 60$
 - Consider a **photon emitted at 90°** w.r.t. the z-axis (beam) in the CM: $(p_{z,CM} = 0, E_{CM}^\gamma = p_T)$
 - $\begin{pmatrix} E_{Lab} \\ p_{z,Lab} \end{pmatrix} = \begin{pmatrix} \gamma & \gamma\beta \\ \gamma\beta & \gamma \end{pmatrix} \begin{pmatrix} p_T \\ 0 \end{pmatrix}$
 - $p_{z,Lab} \simeq 60p_T$! [A 67 MeV γ from a π^0 at rest in the CM can easily be detected.]
- Angle in the Lab. frame: $\tan \theta = \frac{p_T}{p_{z,Lab}} = \frac{1}{\gamma\beta} \Rightarrow \theta \simeq 1^\circ$.
[Rapidity shift: $\Delta y = \tanh^{-1} \beta \simeq 4.8$]
- The entire forward CM hemisphere ($y_{CM} > 0$) within $0^\circ \leq \theta_{Lab} \leq 1^\circ$
[$y_{CM} = 0 \Rightarrow y_{Lab} \simeq 4.8$]
- **Good thing**: small forward detector \equiv large acceptance
- **Bad thing**: high multiplicity \Rightarrow absorber \Rightarrow physics limitation

Backward physics ?

- Let's adopt a **novel strategy** and look at **larger angles**


Backward physics ?

- Let's adopt a **novel strategy** and look at **larger angles**
- Advantages:
 - reduced multiplicities at large(r) angles
 - **access to partons with momentum fraction $x \rightarrow 1$ in the target**
 - last, but not least, the beam pipe is in practice
not a geometrical constrain at $\theta_{CM} \simeq 180^\circ$


Backward physics ?

- Let's adopt a **novel strategy** and look at **larger angles**
- Advantages:
 - reduced multiplicities at large(r) angles
 - **access to partons with momentum fraction $x \rightarrow 1$ in the target**
 - last, but not least, the beam pipe is in practice
not a geometrical constrain at $\theta_{CM} \simeq 180^\circ$

Hadron center-of-mass system



Target rest frame




Backward physics ?

- Let's adopt a **novel strategy** and look at **larger angles**
- Advantages:
 - reduced multiplicities at large(r) angles
 - **access to partons with momentum fraction $x \rightarrow 1$ in the target**
 - last, but not least, the beam pipe is in practice
not a geometrical constrain at $\theta_{CM} \simeq 180^\circ$

Hadron center-of-mass system

Target rest frame




Backward physics ?

- Let's adopt a **novel strategy** and look at **larger angles**
- Advantages:
 - reduced multiplicities at large(r) angles
 - **access to partons with momentum fraction $x \rightarrow 1$ in the target**
 - last, but not least, the beam pipe is in practice
not a geometrical constrain at $\theta_{CM} \simeq 180^\circ$

Hadron center-of-mass system

Target rest frame



backward physics = large- x_2 physics


First systematic access to the target-rapidity region

($x_F \rightarrow -1$)

First systematic access to the target-rapidity region

($x_F \rightarrow -1$)

J/ψ suppression in pA collisions




- x_F systematically studied at fixed target experiments up to +1

First systematic access to the target-rapidity region

($x_F \rightarrow -1$)

J/ψ suppression in pA collisions




- x_F systematically studied at fixed target experiments **up to +1**
- Hera-B was the only one to really explore $x_F < 0$, up to -0.3

First systematic access to the target-rapidity region

($x_F \rightarrow -1$)

J/ψ suppression in pA collisions




- x_F systematically studied at fixed target experiments up to +1
- Hera-B was the only one to really explore $x_F < 0$, up to -0.3
- PHENIX @ RHIC: $-0.1 < x_F < 0.1$ [could be wider with Υ , but low stat.]
- CMS/ATLAS: $|x_F| < 5 \cdot 10^{-3}$; LHCb: $5 \cdot 10^{-3} < x_F < 4 \cdot 10^{-2}$

First systematic access to the target-rapidity region

($x_F \rightarrow -1$)

J/ψ suppression in pA collisions




- x_F systematically studied at fixed target experiments up to +1
- Hera-B was the only one to really explore $x_F < 0$, up to -0.3
- PHENIX @ RHIC: $-0.1 < x_F < 0.1$ [could be wider with Υ , but low stat.]
- CMS/ATLAS: $|x_F| < 5 \cdot 10^{-3}$; LHCb: $5 \cdot 10^{-3} < x_F < 4 \cdot 10^{-2}$

First systematic access to the target-rapidity region

($x_F \rightarrow -1$)

J/ψ suppression in pA collisions



- x_F systematically studied at fixed target experiments **up to +1**
- Hera-B was the only one to really explore $x_F < 0$, up to -0.3
- PHENIX @ RHIC: $-0.1 < x_F < 0.1$ [could be wider with Υ , but low stat.]
- CMS/ATLAS: $|x_F| < 5 \cdot 10^{-3}$; LHCb: $5 \cdot 10^{-3} < x_F < 4 \cdot 10^{-2}$
- If we measure $\Upsilon(b\bar{b})$ at $y_{\text{cms}} \simeq -2.5 \Rightarrow x_F \simeq \frac{2m_\Upsilon}{\sqrt{s}} \sinh(y_{\text{cms}}) \simeq -1$

The lead-ion beam

- Design LHC lead-beam energy: **2.76 TeV** per nucleon

The lead-ion beam

- Design LHC lead-beam energy: **2.76 TeV** per nucleon
- In the fixed target mode, PbA collisions at $\sqrt{s_{NN}} \simeq$ **72 GeV**

The lead-ion beam

- Design LHC lead-beam energy: **2.76 TeV** per nucleon
- In the fixed target mode, PbA collisions at $\sqrt{s_{NN}} \simeq$ **72 GeV**
- Half way **between BNL-RHIC** (AuAu, CuCu @ **200 GeV**) and **CERN-SPS** (PbPb @ **17.2 GeV**)

The lead-ion beam

- Design LHC lead-beam energy: **2.76 TeV** per nucleon
- In the fixed target mode, PbA collisions at $\sqrt{s_{NN}} \simeq \mathbf{72\ TeV}$
- Half way **between BNL-RHIC** (AuAu, CuCu @ **200 GeV**) and **CERN-SPS** (PbPb @ **17.2 GeV**)
- Example of motivations:




Fig. 7. Measured J/ψ production yields, normalised to the yields expected assuming that the only source of suppression is the ordinary absorption by the nuclear medium. The data is shown as a function of the energy density reached in the several collision systems.

The lead-ion beam

- Design LHC lead-beam energy: **2.76 TeV** per nucleon
- In the fixed target mode, PbA collisions at $\sqrt{s_{NN}} \simeq 72$ **GeV**
- Half way **between BNL-RHIC** (AuAu, CuCu @ **200 GeV**) and **CERN-SPS** (PbPb @ **17.2 GeV**)
- Example of motivations:




Fig. 7. Measured J/ψ production yields, normalised to the yields expected assuming that the only source of suppression is the ordinary absorption by the nuclear medium. The data is shown as a function of the energy density reached in the several collision systems.

The beam extraction


- ★ The LHC beam may be extracted using “Strong crystalline field”
without any decrease in performance of the LHC !

E. Uggerhøj, U.I Uggerhøj, NIM B 234 (2005) 31, Rev. Mod. Phys. 77 (2005) 1131

The beam extraction

- ★ The LHC beam may be extracted using “Strong crystalline field”
without any decrease in performance of the LHC !


E. Uggerhøj, U.I Uggerhøj, NIM B 234 (2005) 31, Rev. Mod. Phys. 77 (2005) 1131





The beam extraction

- ★ The LHC beam may be extracted using “Strong crystalline field”
without any decrease in performance of the LHC !

E. Uggerhøj, U.I Uggerhøj, NIM B 234 (2005) 31, Rev. Mod. Phys. 77 (2005) 1131




- ★ **Illustration for collimation**





The beam extraction

- ★ The LHC beam may be extracted using “Strong crystalline field”
without any decrease in performance of the LHC !

E. Uggerhøj, U.I Uggerhøj, NIM B 234 (2005) 31, Rev. Mod. Phys. 77 (2005) 1131





- ★ Illustration for collimation



- ★ Tests will be performed on the LHC beam:
LUA9 proposal approved by the LHCC



The beam extraction

- Inter-crystalline fields are huge



The beam extraction



- Inter-crystalline fields are huge



- The channeling efficiency is high for a deflection of a few mrad

The beam extraction



- Inter-crystalline fields are huge



- The **channeling efficiency** is high for a deflection of a few mrad
- One can **extract** a significant part of the **beam loss** ($10^9 p^+ s^{-1}$)

The beam extraction

- Inter-crystalline fields are huge



- The **channeling efficiency** is high for a deflection of a few mrad
- One can **extract** a significant part of the **beam loss** ($10^9 p^+ s^{-1}$)
- Simple and robust way to extract the most energetic beam ever:



Luminosities

- Expected **proton flux** $\Phi_{beam} = 5 \times 10^8 p^+ s^{-1}$

Luminosities

- Expected **proton flux** $\Phi_{beam} = 5 \times 10^8 \text{ p}^+ \text{ s}^{-1}$
- Instantaneous **Luminosity**:

$$\mathcal{L} = \Phi_{beam} \times N_{target} = N_{beam} \times (\rho \times \ell \times \mathcal{N}_A) / A$$

[ℓ : target thickness (for instance 1cm)]

Luminosities

- Expected **proton flux** $\Phi_{beam} = 5 \times 10^8 \text{ p}^+ \text{s}^{-1}$
- Instantaneous **Luminosity**:

$$\mathcal{L} = \Phi_{beam} \times N_{target} = N_{beam} \times (\rho \times \ell \times \mathcal{N}_A) / A$$

[ℓ : target thickness (for instance 1cm)]

- Integrated luminosity: $\int dt \mathcal{L}$ over **10^7 s for p^+** and **10^6 for Pb**
[the so-called LHC years]

Luminosities

- Expected **proton flux** $\Phi_{beam} = 5 \times 10^8 \text{ p}^+ \text{s}^{-1}$
- Instantaneous **Luminosity**:

$$\mathcal{L} = \Phi_{beam} \times N_{target} = N_{beam} \times (\rho \times \ell \times \mathcal{N}_A) / A$$

[ℓ : target thickness (for instance 1cm)]

- Integrated luminosity: $\int dt \mathcal{L}$ over **10^7 s for p^+** and **10^6 for Pb**

[the so-called LHC years]

Target	$\rho \text{ (g.cm}^{-3}\text{)}$	A	$\mathcal{L} \text{ (}\mu\text{b}^{-1}.\text{s}^{-1}\text{)}$	$\int \mathcal{L} \text{ (pb}^{-1}.\text{yr}^{-1}\text{)}$
Sol. H₂	0.09	1	26	260
Liq. H₂	0.07	1	20	200
Liq. D₂	0.16	2	24	240
Be	1.85	9	62	620
Cu	8.96	64	42	420
W	19.1	185	31	310
Pb	11.35	207	16	160

Luminosities

- 1 meter-long liquid H_2 & D_2 targets can be used (see NA51, ...)


Luminosities

- 1 meter-long liquid H_2 & D_2 targets can be used (see NA51, ...)
- This gives: $\mathcal{L}_{H_2/D_2} \simeq 20 \text{ fb}^{-1} \text{ y}^{-1}$

Luminosities

- 1 meter-long liquid H_2 & D_2 targets can be used (see NA51, ...)
- This gives: $\mathcal{L}_{H_2/D_2} \simeq 20 \text{ fb}^{-1} \text{ y}^{-1}$
- Recycling the LHC beam loss, one gets

a luminosity comparable to the LHC itself !




Luminosities

- 1 meter-long liquid H_2 & D_2 targets can be used (see NA51, ...)
- This gives: $\mathcal{L}_{H_2/D_2} \simeq 20 \text{ fb}^{-1} \text{ y}^{-1}$
- Recycling the LHC beam loss, one gets

a luminosity comparable to the LHC itself !

- PHENIX lumi in their decadal plan
 - Run14pp 12 pb^{-1} @ $\sqrt{s_{NN}} = 200 \text{ GeV}$
 - Run14dAu 0.15 pb^{-1} @ $\sqrt{s_{NN}} = 200 \text{ GeV}$




Luminosities

- 1 meter-long liquid H_2 & D_2 targets can be used (see NA51, ...)
- This gives: $\mathcal{L}_{H_2/D_2} \simeq 20 \text{ fb}^{-1} \text{ y}^{-1}$
- Recycling the LHC beam loss, one gets

a luminosity comparable to the LHC itself !

- PHENIX lumi in their decadal plan
 - Run14pp 12 pb^{-1} @ $\sqrt{s_{NN}} = 200 \text{ GeV}$
 - Run14dAu 0.15 pb^{-1} @ $\sqrt{s_{NN}} = 200 \text{ GeV}$
- AFTER vs PHENIX@RHIC:
3 orders of magnitude larger




Luminosities

- 1 meter-long liquid H_2 & D_2 targets can be used (see NA51, ...)
- This gives: $\mathcal{L}_{H_2/D_2} \simeq 20 \text{ fb}^{-1} \text{ y}^{-1}$
- Recycling the LHC beam loss, one gets

a luminosity comparable to the LHC itself !

- PHENIX lumi in their decadal plan
 - Run14pp 12 pb^{-1} @ $\sqrt{s_{NN}} = 200 \text{ GeV}$
 - Run14dAu 0.15 pb^{-1} @ $\sqrt{s_{NN}} = 200 \text{ GeV}$
- AFTER vs PHENIX@RHIC:
3 orders of magnitude larger
- Lumi for Pb runs in the backup slides
(roughly 10 times that planned for the LHC)



A few figures on the (extracted) proton beam

- Beam loss: $10^9 \text{ p}^+ \text{s}^{-1}$
- Extracted intensity: $5 \times 10^8 \text{ p}^+ \text{s}^{-1}$ (1/2 the beam loss) E. Uggerhøj, U.I Uggerhøj, NIM B 234 (2005) 31

A few figures on the (extracted) proton beam

- Beam loss: $10^9 p^+ s^{-1}$
- Extracted intensity: $5 \times 10^8 p^+ s^{-1}$ (1/2 the beam loss) E. Uggerhøj, U.I Uggerhøj, NIM B 234 (2005) 31
- Number of p^+ : 2808 bunches of $1.15 \times 10^{11} p^+ = 3.2 \times 10^{14} p^+$

A few figures on the (extracted) proton beam

- Beam loss: $10^9 p^+ s^{-1}$
- Extracted intensity: $5 \times 10^8 p^+ s^{-1}$ (1/2 the beam loss) E. Uggerhøj, U.I Uggerhøj, NIM B 234 (2005) 31
- Number of p^+ : 2808 bunches of $1.15 \times 10^{11} p^+ = 3.2 \times 10^{14} p^+$
- Revolution frequency: Each bunch passes the extraction point at a rate of $3.10^5 \text{ km.s}^{-1} / 27 \text{ km} \simeq 11 \text{ kHz}$

A few figures on the (extracted) proton beam

- Beam loss: $10^9 p^+ s^{-1}$
- Extracted intensity: $5 \times 10^8 p^+ s^{-1}$ (1/2 the beam loss) E. Uggerhøj, U.I Uggerhøj, NIM B 234 (2005) 31
- Number of p^+ : 2808 bunches of $1.15 \times 10^{11} p^+ = 3.2 \times 10^{14} p^+$
- Revolution frequency: Each bunch passes the extraction point at a rate of $3.10^5 \text{ km.s}^{-1} / 27 \text{ km} \simeq 11 \text{ kHz}$
- Extracted “mini” bunches:
 - the crystal sees $2808 \times 11000 \text{ s}^{-1} \simeq 3.10^7 \text{ bunches s}^{-1}$
 - one extracts $5.10^8 / 3.10^7 \simeq 15 p^+$ from each bunch at each pass
 - Provided that the probability of interaction with the target is below 5%,

no pile-up !

A few figures on the (extracted) proton beam

- Beam loss: $10^9 p^+ s^{-1}$
- Extracted intensity: $5 \times 10^8 p^+ s^{-1}$ (1/2 the beam loss) E. Uggerhøj, U.I Uggerhøj, NIM B 234 (2005) 31
- Number of p^+ : 2808 bunches of $1.15 \times 10^{11} p^+ = 3.2 \times 10^{14} p^+$
- Revolution frequency: Each bunch passes the extraction point at a rate of $3.10^5 \text{ km.s}^{-1} / 27 \text{ km} \simeq 11 \text{ kHz}$
- Extracted “mini” bunches:
 - the crystal sees $2808 \times 11000 \text{ s}^{-1} \simeq 3.10^7 \text{ bunches s}^{-1}$
 - one extracts $5.10^8 / 3.10^7 \simeq 15 p^+$ from each bunch at each pass
 - Provided that the probability of interaction with the target is below 5%,
- Extraction over a 10h fill:

no pile-up !

 - $5 \times 10^8 p^+ \times 3600 \text{ s h}^{-1} \times 10 \text{ h} = 1.8 \times 10^{13} p^+ \text{ fill}^{-1}$
 - This means $1.8 \times 10^{13} / 3.2 \times 10^{14} \simeq 5.6\%$ of the p^+ in the beam

These protons are lost anyway !

A few figures on the (extracted) proton beam

- Beam loss: $10^9 p^+ s^{-1}$
- Extracted intensity: $5 \times 10^8 p^+ s^{-1}$ (1/2 the beam loss) E. Uggerhøj, U.I Uggerhøj, NIM B 234 (2005) 31
- Number of p^+ : 2808 bunches of $1.15 \times 10^{11} p^+ = 3.2 \times 10^{14} p^+$
- Revolution frequency: Each bunch passes the extraction point at a rate of $3.10^5 \text{ km.s}^{-1} / 27 \text{ km} \simeq 11 \text{ kHz}$
- Extracted “mini” bunches:
 - the crystal sees $2808 \times 11000 \text{ s}^{-1} \simeq 3.10^7 \text{ bunches s}^{-1}$
 - one extracts $5.10^8 / 3.10^7 \simeq 15 p^+$ from each bunch at each pass
 - Provided that the probability of interaction with the target is below 5%,
- Extraction over a 10h fill:

no pile-up !

 - $5 \times 10^8 p^+ \times 3600 \text{ s h}^{-1} \times 10 \text{ h} = 1.8 \times 10^{13} p^+ \text{ fill}^{-1}$
 - This means $1.8 \times 10^{13} / 3.2 \times 10^{14} \simeq 5.6\%$ of the p^+ in the beam
These protons are lost anyway !
- similar figures for the Pb-beam extraction

Part III


AFTER: flagship measurements

Key studies: gluons in the proton

- Gluon distribution at mid, high and ultra-high x_B in the proton


Key studies: gluons in the proton

- **Gluon distribution** at mid, high and ultra-high x_B in the proton
 - Not easily accessible in DIS



Key studies: gluons in the proton


- **Gluon distribution** at mid, high and ultra-high x_B in the proton
 - Not easily accessible in DIS
 - Very large uncertainties



Key studies: gluons in the proton

- **Gluon distribution** at mid, high and ultra-high x_B in the proton
 - Not easily accessible in DIS
 - Very large uncertainties

Accessible thanks gluon sensitive probes,




Key studies: gluons in the proton

- **Gluon distribution** at mid, high and ultra-high x_B in the proton
 - Not easily accessible in DIS
 - Very large uncertainties

Accessible thanks gluon sensitive probes,

- **quarkonia**
see a recent study by D. Diakonov *et al.*, JHEP 1302 (2013) 069



Key studies: gluons in the proton

- **Gluon distribution** at mid, high and ultra-high x_B in the proton
 - Not easily accessible in DIS
 - Very large uncertainties


Accessible thanks gluon sensitive probes,

- **quarkonia**

see a recent study by D. Diakonov *et al.*, JHEP 1302 (2013) 069

- **Isolated photon**

see the recent survey by D. d'Enterria, R. Rojo, Nucl.Phys. B860 (2012) 311



Key studies: gluons in the proton

- **Gluon distribution** at mid, high and ultra-high x_B in the proton
 - Not easily accessible in DIS
 - Very large uncertainties

Accessible thanks gluon sensitive probes,


- **quarkonia**

see a recent study by D. Diakonov *et al.*, JHEP 1302 (2013) 069

- **Isolated photon**

see the recent survey by D. d'Enterria, R. Rojo, Nucl.Phys. B860 (2012) 311

- **jets** ($P_T \in [20, 40]$ GeV)



Key studies: gluons in the proton

- **Gluon distribution** at mid, high and ultra-high x_B in the proton
 - Not easily accessible in DIS
 - Very large uncertainties

Accessible thanks gluon sensitive probes,

- **quarkonia**


see a recent study by D. Diakonov *et al.*, JHEP 1302 (2013) 069

- **Isolated photon**

see the recent survey by D. d'Enterria, R. Rojo, Nucl.Phys. B860 (2012) 311

- **jets** ($P_T \in [20, 40]$ GeV)

Multiple probes needed to **check factorisation**



Isolated- γ in p(7 TeV)-p(rest): $\sqrt{s} \sim 115$ GeV

- p-p photon kinematics at fixed-target LHC (central rapidities):
To access $x > 0.3$ one needs isolated- γ at: $p_T = x_T \sqrt{s}/2 > 20$ GeV/c


- JETPHOX NLO
pQCD calculations:

p-p at $\sqrt{s}=115$ GeV

$|y| < 0.5$, $p_T > 20$ GeV/c

Isolation: $R=0.4$, $E_T^{\text{had}} < 5$ GeV

\mathcal{L} (10 cm H_2 -target) $\sim 2 \cdot 10^3$ pb $^{-1}$ /year




PDF: CT10 52 eigenval. (90% CL)

Scales: $\mu_i = p_T$


FF = BFG-II

x-section uncertainties^(*) of $\pm 150\%$

^(*) (68%CL)/(90% CL) ~ 1.65




Key studies: gluons in the neutron



Gluon PDF for the neutron unknown

Key studies: gluons in the neutron




Gluon PDF for the neutron unknown

possible experimental probes

- heavy quarkonia
- isolated photons
- jets


Key studies: gluons in the neutron



Gluon PDF for the neutron unknown

possible experimental probes


- heavy quarkonia
- isolated photons
- jets



Pioneer measurement by E866

- using $\Upsilon \rightarrow Q^2 \simeq 100 \text{ GeV}^2$
- outcome: $g_n(x) \simeq g_p(x)$


Key studies: gluons in the neutron



Gluon PDF for the neutron unknown

possible experimental probes

- heavy quarkonia
- isolated photons
- jets




Pioneer measurement by E866

- using $\Upsilon \rightarrow Q^2 \simeq 100 \text{ GeV}^2$
- outcome: $g_n(x) \simeq g_p(x)$

could be extended with AFTER

- using J/ψ , ..., $C = +1$ onia, ...
- wider x range & lower Q^2


Key studies: gluons in the neutron



Gluon PDF for the neutron unknown

possible experimental probes

- heavy quarkonia
- isolated photons
- jets



Pioneer measurement by E866

- using $\Upsilon \rightarrow Q^2 \simeq 100 \text{ GeV}^2$
- outcome: $g_n(x) \simeq g_p(x)$

could be extended with AFTER

- using J/ψ , ..., $C = +1$ onia, ...
- wider x range & lower Q^2


target	yearly lumi	$\mathcal{B} \frac{dN_{J/\psi}}{dy}$	$\mathcal{B} \frac{dN_r}{dy}$
1m Liq. H ₂	20 fb ⁻¹	4.0×10^8	9.0×10^5
1m Liq. D ₂	24 fb ⁻¹	9.6×10^8	1.9×10^6

Key studies

- Heavy-quark distributions (at high x_B)


Key studies

- Heavy-quark distributions (at high x_B)
 - Pin down intrinsic charm, ... at last




Key studies

- Heavy-quark distributions (at high x_B)
 - Pin down **intrinsic charm**, ... at last
 - **Total open charm and beauty** cross section (aim: down to $P_T \rightarrow 0$)




Key studies

- Heavy-quark distributions (at high x_B)
 - Pin down **intrinsic charm**, ... at last
 - **Total open charm and beauty** cross section
(aim: down to $P_T \rightarrow 0$)
requires



Key studies

- Heavy-quark distributions (at high x_B)
 - Pin down **intrinsic charm**, ... at last
 - **Total open charm and beauty** cross section (aim: down to $P_T \rightarrow 0$)
- requires
- several **complementary** measurements




Key studies

- Heavy-quark distributions (at high x_B)
 - Pin down **intrinsic charm**, ... at last
 - **Total open charm and beauty** cross section (aim: down to $P_T \rightarrow 0$)

requires

- several **complementary** measurements
- good coverage in the **target-rapidity region**




Key studies

- **Heavy-quark** distributions (at high x_B)
 - Pin down **intrinsic charm**, ... at last
 - **Total open charm and beauty** cross section (aim: down to $P_T \rightarrow 0$)

requires

- several **complementary** measurements
- good coverage in the **target-rapidity** region
- high **luminosity** to reach large x_B







Key studies

- **Heavy-quark** distributions (at high x_B)
 - Pin down **intrinsic charm**, ... at last
 - **Total open charm and beauty** cross section (aim: down to $P_T \rightarrow 0$)

requires

- several **complementary** measurements
- good coverage in the **target-rapidity region**
- high **luminosity** to reach large x_B



3 sets from CTEQ6c (Pumplin *et al.*)

Key studies: gluon contribution to the proton spin

- Gluon Sivers effect: correlation between the gluon transverse momentum & the proton spin

Key studies: gluon contribution to the proton spin



- **Gluon Sivers effect**: correlation between the **gluon transverse momentum** & the **proton spin**
 - Transverse **single spin asymmetries** using **gluon sensitive probes**

Key studies: gluon contribution to the proton spin



- **Gluon Sivers effect**: correlation between the **gluon transverse momentum** & the **proton spin**
 - Transverse **single spin asymmetries** using **gluon sensitive probes**
 - quarkonia (J/ψ , Υ , χ_c , ...)

F. Yuan, PRD 78 (2008) 014024

Key studies: gluon contribution to the proton spin



- **Gluon Sivers effect**: correlation between the **gluon transverse momentum** & the **proton spin**
 - Transverse **single spin asymmetries** using **gluon sensitive probes**
- quarkonia (J/ψ , Υ , χ_c , ...)
- B & D meson production

F. Yuan, PRD 78 (2008) 014024

Key studies: gluon contribution to the proton spin



- **Gluon Sivers effect:** correlation between the **gluon transverse momentum** & the **proton spin**
 - Transverse **single spin asymmetries** using **gluon sensitive probes**
 - quarkonia (J/ψ , Υ , χ_c , ...)
 - B & D meson production
 - γ , γ -jet, $\gamma - \gamma$

F. Yuan, PRD 78 (2008) 014024

A. Bacchetta, *et al.*, PRL 99 (2007) 212002
 J.W. Qiu, *et al.*, PRL 107 (2011) 062001

Key studies: gluon contribution to the proton spin



- **Gluon Sivers effect:** correlation between the **gluon transverse momentum** & the **proton spin**
 - Transverse **single spin asymmetries** using **gluon sensitive probes**
- quarkonia (J/ψ , Υ , χ_c , ...)
- B & D meson production
- γ , γ -jet, $\gamma - \gamma$
- the target-rapidity region corresponds to **high x^\uparrow** where the **k_T -spin correlation is the largest**

F. Yuan, PRD 78 (2008) 014024

A. Bacchetta, *et al.*, PRL 99 (2007) 212002
J.W. Qiu, *et al.*, PRL 107 (2011) 062001

Key studies: gluon contribution to the proton spin



- **Gluon Sivers effect**: correlation between the **gluon transverse momentum** & the **proton spin**
 - Transverse **single spin asymmetries** using **gluon sensitive probes**
- quarkonia (J/ψ , Υ , χ_c , ...)
- B & D meson production
- γ , γ -jet, $\gamma - \gamma$
- the target-rapidity region corresponds to **high x^\uparrow** where the **k_T -spin correlation is the largest**
- In general, one can carry out an extensive spin-physics program

F. Yuan, PRD 78 (2008) 014024

A. Bacchetta, *et al.*, PRL 99 (2007) 212002
 J.W. Qiu, *et al.*, PRL 107 (2011) 062001

Key studies: gluon contribution to the proton spin



- **Gluon Sivers effect: correlation** between the **gluon transverse momentum** & the **proton spin**

- Transverse

expected Sivers asymmetry in
D-Y@AFTER, sign change,
no TMD evolution

- quark

e probes

- B & C

3 (2008) 014024

- γ , γ -jet


12002
101

- the target

e largest

- In general

rogram



M. Anselmino, Trento, February 2013

Key studies: gluon contribution to the proton spin



- **Gluon Sivers effect:** correlation between the **gluon transverse momentum** & the **proton spin**
- Transverse **single spin asymmetries** using **gluon sensitive probes**
- quarkonia (J/ψ , Υ , χ_c , ...)

F. Yuan, PRD 78 (2008) 014024

PHYSICAL REVIEW D 86, 094007 (2012)

Polarized gluon studies with charmonium and bottomonium at LHCb and AFTER

Daniël Boer*


*Theory Group, KVI, University of Groningen, Zernikelaan 25, NL-9747 AA Groningen, The Netherlands*Cristian Pisano[†]*Istituto Nazionale di Fisica Nucleare, Sezione di Cagliari, C.P. 170, I-09042 Monserrato (CA), Italy*

- In general, one can carry out an extensive spin-physics program

Key studies: large- x gluon content of the nucleus


Key studies: large-x gluon content of the nucleus

- Large-x gluon nPDF: unknown
- Gluon EMC effect ?




Key studies: large- x gluon content of the nucleus

- Large- x gluon nPDF: unknown
- Gluon EMC effect ?
- Hint from Υ data at RHIC






Key studies: large-x gluon content of the nucleus

- Large-x gluon nPDF: unknown
- Gluon EMC effect ?
- Hint from Υ data at RHIC
- Strongly limited in terms of statistics
after 10 years of RHIC:






Key studies: large-x gluon content of the nucleus

- Large-x gluon nPDF: unknown
- Gluon EMC effect ?
- Hint from Υ data at RHIC
- Strongly limited in terms of statistics
after 10 years of RHIC:
- DIS contribution expected for low x mainly
projected contribution of LHeC:



Key studies: large- x gluon content of the nucleus

- Large- x gluon nPDF: unknown
- Gluon EMC effect ?
- Hint from Υ data at RHIC
- Strongly limited in terms of statistics
after 10 years of RHIC:
- DIS contribution expected for low x mainly
projected contribution of LHeC:
- AFTER allows for extensive studies of
gluon sensitive probes in pA
- Unique potential for gluons at $x > 0.1$



Precision heavy-flavour studies in Heavy-Ion Collisions


- Very precise pp and pA baselines (yields, A & y dependences)

Precision heavy-flavour studies in Heavy-Ion Collisions

- Very precise pp and pA baselines (yields, A & y dependences)
- Modern technologies to look for quarkonium excited states

Precision heavy-flavour studies in Heavy-Ion Collisions


- Very precise *pp* and *pA* **baselines** (yields, A & y dependences)
- **Modern technologies** to look for quarkonium **excited states**



HERA-B PRD 79 (2009)
012001, and ref. therein

Precision heavy-flavour studies in Heavy-Ion Collisions


- Very precise *pp* and *pA* **baselines** (yields, A & y dependences)
- **Modern technologies** to look for quarkonium **excited states**
- Energy between SPS and RHIC:
QGP should be formed **w/o $c\bar{c}$ recombination**



HERA-B PRD 79 (2009)
012001, and ref. therein

Precision heavy-flavour studies in Heavy-Ion Collisions


- Very precise *pp* and *pA* **baselines** (yields, A & y dependences)
- **Modern technologies** to look for quarkonium **excited states**
- Energy between SPS and RHIC:
QGP should be formed **w/o $c\bar{c}$ recombination**
- **Open heavy-flavour measurement**
down to $P_T = 0$ thanks to the boost.



HERA-B PRD 79 (2009)
012001, and ref. therein

Precision heavy-flavour studies in Heavy-Ion Collisions

- Very precise *pp* and *pA* **baselines** (yields, A & y dependences)
- **Modern technologies** to look for quarkonium **excited states**
- Energy between SPS and RHIC:
QGP should be formed **w/o $c\bar{c}$ recombination**
- **Open heavy-flavour measurement**
down to $P_T = 0$ thanks to the boost.
- Real hope of being able to look at the **quarkonium sequential suppression**



HERA-B PRD 79 (2009)
012001, and ref. therein

Key studies: W/Z production at threshold

- For the first time, one would study W/Z production in their threshold region ($m_{W/Z}/\sqrt{s_{AFTER}} \sim 1$)

Key studies: W/Z production at threshold

- For the first time, one would study W/Z production in their **threshold region** ($m_{W/Z}/\sqrt{s_{AFTER}} \sim 1$)
 - Unique opportunity to measure QCD/threshold effects on W/Z production

Key studies: W/Z production at threshold

- For the first time, one would study W/Z production in their **threshold region** ($m_{W/Z}/\sqrt{s_{AFTER}} \sim 1$)
 - Unique opportunity to measure QCD/threshold effects on W/Z production
 - If W'/Z' exist, their production may share similar threshold corrections to that of W/Z , but at LHC energies ($m_{W'/Z'}/\sqrt{s_{LHC}} \sim 1$?)

Key studies: W/Z production at threshold

- For the first time, one would study W/Z production in their **threshold region** ($m_{W/Z}/\sqrt{s_{AFTER}} \sim 1$)
 - Unique opportunity to measure QCD/threshold effects on W/Z production
- If W'/Z' exist, their production may share similar threshold corrections to that of W/Z , but at LHC energies ($m_{W'/Z'}/\sqrt{s_{LHC}} \sim 1$?)
- Reconstructed rate are most likely between **a few dozen to a few thousand / year**

Further key studies ?

(Multiply) heavy baryons:

Further key studies ?

(Multiply) heavy baryons:

- $\Lambda_b \rightarrow \Lambda J/\psi$

Further key studies ?

(Multiply) heavy baryons:

- $\Lambda_b \rightarrow \Lambda J/\psi$
 - $d\sigma(b)/dy|_{y=0} \gtrsim 100 \text{ nb}$

Further key studies ?

(Multiply) heavy baryons:

- $\Lambda_b \rightarrow \Lambda J/\psi$
 - $d\sigma(b)/dy|_{y=0} \gtrsim 100 \text{ nb}$
 - $\mathcal{N}(b)/\text{year} \simeq 2 \times 100 \times 10^6 \times 20 = 4 \times 10^9$

Further key studies ?

(Multiply) heavy baryons:

- $\Lambda_b \rightarrow \Lambda J/\psi$
 - $d\sigma(b)/dy|_{y=0} \gtrsim 100 \text{ nb}$
 - $\mathcal{N}(b)/\text{year} \simeq 2 \times 100 \times 10^6 \times 20 = 4 \times 10^9$
 - $\mathcal{B}(b \rightarrow \Lambda_b) \times \mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda) = 5.8 \pm 0.8 \times 10^{-5}$
 $(\mathcal{B}(J/\psi \rightarrow \mu\mu) = 6\%)$

Further key studies ?

(Multiply) heavy baryons:

- $\Lambda_b \rightarrow \Lambda J/\psi$
 - $d\sigma(b)/dy|_{y=0} \gtrsim 100 \text{ nb}$
 - $\mathcal{N}(b)/\text{year} \simeq 2 \times 100 \times 10^6 \times 20 = 4 \times 10^9$
 - $\mathcal{B}(b \rightarrow \Lambda_b) \times \mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda) = 5.8 \pm 0.8 \times 10^{-5}$
 $(\mathcal{B}(J/\psi \rightarrow \mu\mu) = 6\%)$
 - 15 000 $\Lambda_b \rightarrow J/\psi \Lambda \rightarrow \mu^+ \mu^- \Lambda$ events: enough to perform a polarisation measurement

see e.g. LHCb arXiv:1302.5578 [hep-ex]

Further key studies ?

(Multiply) heavy baryons:

- $\Lambda_b \rightarrow \Lambda J/\psi$
 - $d\sigma(b)/dy|_{y=0} \gtrsim 100 \text{ nb}$
 - $\mathcal{N}(b)/\text{year} \simeq 2 \times 100 \times 10^6 \times 20 = 4 \times 10^9$
 - $\mathcal{B}(b \rightarrow \Lambda_b) \times \mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda) = 5.8 \pm 0.8 \times 10^{-5}$
 $(\mathcal{B}(J/\psi \rightarrow \mu\mu) = 6\%)$
 - 15 000 $\Lambda_b \rightarrow J/\psi \Lambda \rightarrow \mu^+ \mu^- \Lambda$ events: enough to perform a polarisation measurement see e.g. LHCb arXiv:1302.5578 [hep-ex]
- discovery potential ? ($\Xi_{cc}, \Omega^{++}(ccc), \dots$)

Further key studies ?

(Multiply) heavy baryons:

- $\Lambda_b \rightarrow \Lambda J/\psi$
 - $d\sigma(b)/dy|_{y=0} \gtrsim 100 \text{ nb}$
 - $\mathcal{N}(b)/\text{year} \simeq 2 \times 100 \times 10^6 \times 20 = 4 \times 10^9$
 - $\mathcal{B}(b \rightarrow \Lambda_b) \times \mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda) = 5.8 \pm 0.8 \times 10^{-5}$
($\mathcal{B}(J/\psi \rightarrow \mu\mu) = 6\%$)
 - 15 000 $\Lambda_b \rightarrow J/\psi \Lambda \rightarrow \mu^+ \mu^- \Lambda$ events: enough to perform a polarisation measurement see e.g. LHCb arXiv:1302.5578 [hep-ex]
- discovery potential ? ($\Xi_{cc}, \Omega^{++}(ccc), \dots$)
 - Ξ_{cc}, \dots , cross sections in the central region are being calculated with the MC generator GENXICC

C.H. Chang, J.X. Wang, X.G. Wu. Comput.Phys.Commun. 177 (2007) 467

Further key studies ?

(Multiply) heavy baryons:

- $\Lambda_b \rightarrow \Lambda J/\psi$
 - $d\sigma(b)/dy|_{y=0} \gtrsim 100 \text{ nb}$
 - $\mathcal{N}(b)/\text{year} \simeq 2 \times 100 \times 10^6 \times 20 = 4 \times 10^9$
 - $\mathcal{B}(b \rightarrow \Lambda_b) \times \mathcal{B}(\Lambda_b \rightarrow J/\psi \Lambda) = 5.8 \pm 0.8 \times 10^{-5}$
($\mathcal{B}(J/\psi \rightarrow \mu\mu) = 6\%$)
 - 15 000 $\Lambda_b \rightarrow J/\psi \Lambda \rightarrow \mu^+ \mu^- \Lambda$ events: enough to perform a polarisation measurement see e.g. LHCb arXiv:1302.5578 [hep-ex]
- discovery potential ? ($\Xi_{cc}, \Omega^{++}(ccc), \dots$)
 - Ξ_{cc}, \dots , cross sections in the central region are being calculated with the MC generator GENXICC
- they should also be calculated for $x_F \rightarrow -1$

C.H. Chang, J.X. Wang, X.G. Wu. Comput.Phys.Commun. 177 (2007) 467

where IQ could dominate

Further key studies ?

- $\gamma + p$ interaction via **ultra-peripheral collisions**

Further key studies ?

- $\gamma + p$ interaction via **ultra-peripheral collisions**
 - $\gamma_{\text{lab}}^{\text{beam}} \simeq 7000$ ($E_p = 7000$ GeV)
 - $E_{\gamma,\text{lab}}^{\text{max}} \simeq \gamma_{\text{lab}}^{\text{beam}} \times 30$ MeV ($1/R_{\text{Pb}} \simeq 30$ MeV)
 - $\sqrt{s_{\gamma p}} = \sqrt{2m_p E_\gamma}$ up to 20 GeV
 - No pile-up

Further key studies ?

- $\gamma + p$ interaction via **ultra-peripheral collisions**
 - $\gamma_{\text{lab}}^{\text{beam}} \simeq 7000$ ($E_p = 7000$ GeV)
 - $E_{\gamma,\text{lab}}^{\text{max}} \simeq \gamma_{\text{lab}}^{\text{beam}} \times 30$ MeV ($1/R_{\text{Pb}} \simeq 30$ MeV)
 - $\sqrt{s_{\gamma p}} = \sqrt{2m_p E_\gamma}$ up to 20 GeV
 - No pile-up
- **Fracture functions**


Further key studies ?

- $\gamma + p$ interaction via **ultra-peripheral collisions**

- $\gamma_{\text{lab}}^{\text{beam}} \simeq 7000$ ($E_p = 7000$ GeV)
- $E_{\gamma,\text{lab}}^{\text{max}} \simeq \gamma_{\text{lab}}^{\text{beam}} \times 30$ MeV ($1/R_{\text{Pb}} \simeq 30$ MeV)
- $\sqrt{s_{\gamma p}} = \sqrt{2m_p E_\gamma}$ up to 20 GeV
- No pile-up

- **Fracture functions**

- via Drell-Yan pair production
+ identified hadron



L. Trentadue, G. Veneziano, PLB 323 (1994) 201
F. Ceccopieri, L. Trentadue, PLB 668 (2008) 319
talk on Tuesday by O. Samoylov


Further key studies ?

- $\gamma + p$ interaction via **ultra-peripheral collisions**

- $\gamma_{\text{lab}}^{\text{beam}} \simeq 7000$ ($E_p = 7000$ GeV)
- $E_{\gamma,\text{lab}}^{\text{max}} \simeq \gamma_{\text{lab}}^{\text{beam}} \times 30$ MeV ($1/R_{\text{Pb}} \simeq 30$ MeV)
- $\sqrt{s_{\gamma p}} = \sqrt{2m_p E_\gamma}$ up to 20 GeV
- No pile-up

- **Fracture functions**

- via Drell-Yan pair production
+ identified hadron



L. Trentadue, G. Veneziano, PLB 323 (1994) 201
F. Ceccopieri, L. Trentadue, PLB 668 (2008) 319
talk on Tuesday by O. Samoylov

- privileged region for the identified hadron: either the projectile- or **target-rapidity region**


Further key studies ?

- $\gamma + p$ interaction via **ultra-peripheral collisions**

- $\gamma_{\text{lab}}^{\text{beam}} \simeq 7000$ ($E_p = 7000$ GeV)
- $E_{\gamma,\text{lab}}^{\text{max}} \simeq \gamma_{\text{lab}}^{\text{beam}} \times 30$ MeV ($1/R_{\text{Pb}} \simeq 30$ MeV)
- $\sqrt{s_{\gamma p}} = \sqrt{2m_p E_\gamma}$ up to 20 GeV
- No pile-up

- **Fracture functions**

- via Drell-Yan pair production
+ identified hadron



L. Trentadue, G. Veneziano, PLB 323 (1994) 201
F. Ceccopieri, L. Trentadue, PLB 668 (2008) 319
talk on Tuesday by O. Samoylov

- privileged region for the identified hadron: either the projectile- or **target-rapidity region**
- the fixed-target mode is ideal for such studies


Further key studies ?

- $\gamma + p$ interaction via **ultra-peripheral collisions**

- $\gamma_{\text{lab}}^{\text{beam}} \simeq 7000$ ($E_p = 7000$ GeV)
- $E_{\gamma,\text{lab}}^{\text{max}} \simeq \gamma_{\text{lab}}^{\text{beam}} \times 30$ MeV ($1/R_{\text{Pb}} \simeq 30$ MeV)
- $\sqrt{s_{\gamma p}} = \sqrt{2m_p E_\gamma}$ up to 20 GeV
- No pile-up

- **Fracture functions**

- via Drell-Yan pair production
+ identified hadron




L. Trentadue, G. Veneziano, PLB 323 (1994) 201
F. Ceccopieri, L. Trentadue, PLB 668 (2008) 319
talk on Tuesday by O. Samoylov

- privileged region for the identified hadron: either the projectile- or **target-rapidity region**
- the fixed-target mode is ideal for such studies
- good prospects for gluon fracture-function studies !


Overall


Fixed Target @ LHC



Overall


Fixed Target @ LHC






Overall

Fixed Target @ LHC



Overall

Fixed Target @ LHC



More details in

Physics Reports 522 (2013) 239–255



Contents lists available at SciVerse ScienceDirect

Physics Reports

journal homepage: www.elsevier.com/locate/physrep



Physics opportunities of a fixed-target experiment using LHC beams

S.J. Brodsky^a, F. Fleuret^b, C. Hadjidakis^c, J.P. Lansberg^{c,*}

^a SLAC National Accelerator Laboratory, Stanford University, Menlo Park, CA 94025, USA

^b Laboratoire Leprince Ringuet, Ecole polytechnique, CNRS/IN2P3, 91128 Palaiseau, France

^c IPNO, Université Paris-Sud, CNRS/IN2P3, 91406 Orsay, France

Contents

1. Introduction.....	6. Deconfinement in heavy-ion collisions.....
2. Key numbers and features.....	6.1. Quarkonium studies.....
3. Nucleon partonic structure.....	6.2. Jet quenching.....
3.1. Drell–Yan.....	6.3. Direct photon.....
3.2. Gluons in the proton at large x	6.4. Deconfinement and the target rest frame.....
3.2.1. Quarkonia.....	6.5. Nuclear-matter baseline.....
3.2.2. Jets.....	7. W and Z boson production in pp , pd and pA collisions.....
3.2.3. Direct/isolated photons.....	7.1. First measurements in pA
3.3. Gluons in the deuteron and in the neutron.....	7.2. W/Z production in pp and pd
3.4. Charm and bottom in the proton.....	8. Exclusive, semi-exclusive and backward reactions.....
3.4.1. Open-charm production.....	8.1. Ultra-peripheral collisions.....
3.4.2. $J/\psi + D$ meson production.....	8.2. Hard diffractive reactions.....
3.4.3. Heavy-quark plus photon production.....	8.3. Heavy-hadron (diffractive) production at $x_F \rightarrow -1$
4. Spin physics.....	8.4. Very backward physics.....
4.1. Transverse SSA and DY.....	8.5. Direct hadron production.....
4.2. Quarkonium and heavy-quark transverse SSA.....	9. Further potentialities of a high-energy fixed-target set-up.....
4.3. Transverse SSA and photon.....	9.1. D and B physics.....
4.4. Spin asymmetries with a final state polarization.....	9.2. Secondary beams.....
5. Nuclear matter.....	9.3. Forward studies in relation with cosmic shower.....
5.1. Quark nPDF: Drell–Yan in pA and $PbPb$	10. Conclusions.....
5.2. Gluon nPDF.....	Acknowledgments.....
5.2.1. Isolated photons and photon–jet correlations.....	References.....
5.2.2. Precision quarkonium and heavy-flavour studies.....	
5.3. Color filtering, energy loss, Sudakov suppression and hadron break-up in the nucleus.....	

Part IV

Conclusion and outlooks

Conclusion

- Both p and Pb LHC beams can be extracted without disturbing the other experiments

Conclusion

- Both p and Pb LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam $\rightarrow 5 \times 10^8$ protons per sec

Conclusion

- Both p and Pb LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam $\rightarrow 5 \times 10^8$ protons per sec
- This allows for high luminosity pp , pA and PbA collisions at $\sqrt{s} = 115$ GeV and $\sqrt{s_{NN}} = 72$ GeV

Conclusion

- Both p and Pb LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam $\rightarrow 5 \times 10^8$ protons per sec
- This allows for high luminosity pp , pA and PbA collisions at $\sqrt{s} = 115$ GeV and $\sqrt{s_{NN}} = 72$ GeV
- **Example:** precision quarkonium studies taking advantage of

Conclusion

- Both p and Pb LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam $\rightarrow 5 \times 10^8$ protons per sec
- This allows for high luminosity pp , pA and PbA collisions at $\sqrt{s} = 115$ GeV and $\sqrt{s_{NN}} = 72$ GeV
- **Example:** precision quarkonium studies taking advantage of
 - high luminosity (reach in y , P_T , small BR channels)
 - target versatility (nuclear effects, strongly limited at colliders)
 - modern detection techniques (e.g. γ detection with high multiplicity)

Conclusion

- Both p and Pb LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam $\rightarrow 5 \times 10^8$ protons per sec
- This allows for high luminosity pp , pA and PbA collisions at $\sqrt{s} = 115$ GeV and $\sqrt{s_{NN}} = 72$ GeV
- **Example:** precision quarkonium studies taking advantage of
 - high luminosity (reach in y , P_T , small BR channels)
 - target versatility (nuclear effects, strongly limited at colliders)
 - modern detection techniques (e.g. γ detection with high multiplicity)
- This would likely prepare the ground for $g(x, Q^2)$ extraction

Conclusion

- Both p and Pb LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam $\rightarrow 5 \times 10^8$ protons per sec
- This allows for high luminosity pp , pA and PbA collisions at $\sqrt{s} = 115$ GeV and $\sqrt{s_{NN}} = 72$ GeV
- **Example:** precision quarkonium studies taking advantage of
 - high luminosity (reach in y , P_T , small BR channels)
 - target versatility (nuclear effects, strongly limited at colliders)
 - modern detection techniques (e.g. γ detection with high multiplicity)
- This would likely prepare the ground for $g(x, Q^2)$ extraction
- A wealth of possible measurements:
DY, Open b/c , jet correlation, UPC... (not mentioning secondary beams)

Conclusion

- Both p and Pb LHC beams can be extracted without disturbing the other experiments
- Extracting **a few per cent of the beam** $\rightarrow 5 \times 10^8$ **protons per sec**
- This allows for **high luminosity** pp , pA and PbA collisions at $\sqrt{s} = 115$ GeV and $\sqrt{s_{NN}} = 72$ GeV
- **Example:** **precision quarkonium studies** taking advantage of
 - high luminosity (reach in y , P_T , small BR channels)
 - target versatility (nuclear effects, strongly limited at colliders)
 - modern detection techniques (e.g. γ detection with high multiplicity)
- This would likely prepare the ground for **$g(x, Q^2)$ extraction**
- A wealth of possible measurements:
DY, Open b/c , jet correlation, UPC... (not mentioning secondary beams)
- Planned LHC long shutdown (< 2020 ?) could be used to install the extraction system

Conclusion

- Both p and Pb LHC beams can be extracted without disturbing the other experiments
- Extracting **a few per cent of the beam** $\rightarrow 5 \times 10^8$ **protons per sec**
- This allows for **high luminosity** pp , pA and PbA collisions at $\sqrt{s} = 115$ GeV and $\sqrt{s_{NN}} = 72$ GeV
- **Example:** **precision quarkonium studies** taking advantage of
 - high luminosity (reach in y , P_T , small BR channels)
 - target versatility (nuclear effects, strongly limited at colliders)
 - modern detection techniques (e.g. γ detection with high multiplicity)
- This would likely prepare the ground for **$g(x, Q^2)$ extraction**
- A wealth of possible measurements:
DY, Open b/c , jet correlation, UPC... (not mentioning secondary beams)
- Planned LHC long shutdown (< 2020 ?) could be used to install the extraction system
- Very good **complementarity** with electron-ion programs

Outlooks

- First physics paper **Physics Reports 522 (2013) 239**

Outlooks

- First physics paper **Physics Reports 522 (2013) 239**
- A 10-day exploratory workshop at ECT* Trento, February 4-13, 2013
slides at <http://indico.in2p3.fr/event/AFTER@ECTstar>

Outlooks

- First physics paper **Physics Reports 522 (2013) 239**
- A 10-day exploratory workshop at ECT* Trento, February 4-13, 2013
slides at <http://indico.in2p3.fr/event/AFTER@ECTstar>
- NEW: Workshop in **Les Houches on 12-17 January 2014**
<http://indico.in2p3.fr/event/AFTER@LesHouches>

Outlooks

- First physics paper **Physics Reports 522 (2013) 239**
- A 10-day exploratory workshop at ECT* Trento, February 4-13, 2013
slides at <http://indico.in2p3.fr/event/AFTER@ECTstar>
- NEW: Workshop in **Les Houches on 12-17 January 2014**
<http://indico.in2p3.fr/event/AFTER@LesHouches>
- We are looking for **more partners** to

Outlooks

- First physics paper **Physics Reports 522 (2013) 239**
- A 10-day exploratory workshop at ECT* Trento, February 4-13, 2013
slides at <http://indico.in2p3.fr/event/AFTER@ECTstar>
- NEW: Workshop in **Les Houches on 12-17 January 2014**
<http://indico.in2p3.fr/event/AFTER@LesHouches>
- We are looking for **more partners** to
 - do first **simulations** (we are getting ready for fast simulations)

Outlooks

- First physics paper **Physics Reports 522 (2013) 239**
- A 10-day exploratory workshop at ECT* Trento, February 4-13, 2013
slides at <http://indico.in2p3.fr/event/AFTER@ECTstar>
- NEW: Workshop in **Les Houches on 12-17 January 2014**
<http://indico.in2p3.fr/event/AFTER@LesHouches>
- We are looking for **more partners** to
 - do first **simulations** (we are getting ready for fast simulations)
 - think about **possible designs**

Outlooks

- First physics paper **Physics Reports 522 (2013) 239**
- A 10-day exploratory workshop at ECT* Trento, February 4-13, 2013
slides at <http://indico.in2p3.fr/event/AFTER@ECTstar>
- NEW: Workshop in **Les Houches on 12-17 January 2014**
<http://indico.in2p3.fr/event/AFTER@LesHouches>
- We are looking for **more partners** to
 - do first **simulations** (we are getting ready for fast simulations)
 - think about **possible designs**
 - think about the optimal **detector technologies**
 - enlarge the physics case (cosmic rays, flavour physics, ...)

Outlooks

- First physics paper **Physics Reports 522 (2013) 239**
- A 10-day exploratory workshop at ECT* Trento, February 4-13, 2013
slides at <http://indico.in2p3.fr/event/AFTER@ECTstar>
- NEW: Workshop in **Les Houches on 12-17 January 2014**
<http://indico.in2p3.fr/event/AFTER@LesHouches>
- We are looking for **more partners** to
 - do first **simulations** (we are getting ready for fast simulations)
 - think about **possible designs**
 - think about the optimal **detector technologies**
 - enlarge the physics case (cosmic rays, flavour physics, ...)
- **Theorist colleagues** are encouraged to think about **additional ideas of physics**

already 2 papers on the physics at AFTER:

T. Liu, B.Q. Ma, EPJC (2012) 72:2037

D. Boer, C. Pisano, Phys.Rev. D86 (2012) 094007

Outlooks

- First physics paper **Physics Reports 522 (2013) 239**
- A 10-day exploratory workshop at ECT* Trento, February 4-13, 2013
slides at <http://indico.in2p3.fr/event/AFTER@ECTstar>
- NEW: Workshop in **Les Houches on 12-17 January 2014**
<http://indico.in2p3.fr/event/AFTER@LesHouches>
- We are looking for **more partners** to
 - do first **simulations** (we are getting ready for fast simulations)
 - think about **possible designs**
 - think about the optimal **detector technologies**
 - enlarge the physics case (cosmic rays, flavour physics, ...)
- **Theorist colleagues** are encouraged to think about **additional ideas of physics**

already 2 papers on the physics at AFTER:
T. Liu, B.Q. Ma, EPJC (2012) 72:2037
D. Boer, C. Pisano, Phys.Rev. D86 (2012) 094007
- Do not hesitate to contact us

Outlooks

- First physics paper **Physics Reports 522 (2013) 239**
- A 10-day exploratory workshop at ECT* Trento, February 4-13, 2013
slides at <http://indico.in2p3.fr/event/AFTER@ECTstar>
- NEW: Workshop in **Les Houches on 12-17 January 2014**
<http://indico.in2p3.fr/event/AFTER@LesHouches>
- We are looking for **more partners** to
 - do first **simulations** (we are getting ready for fast simulations)
 - think about **possible designs**
 - think about the optimal **detector technologies**
 - enlarge the physics case (cosmic rays, flavour physics, ...)
- **Theorist colleagues** are encouraged to think about **additional ideas of physics**

already 2 papers on the physics at AFTER:
T. Liu, B.Q. Ma, EPJC (2012) 72:2037
D. Boer, C. Pisano, Phys.Rev. D86 (2012) 094007
- Do not hesitate to contact us
- Webpage: <http://after.in2p3.fr>



Part V

Backup slides


Beam extraction

Beam extraction @ LHC

... there are extremely promising possibilities to extract 7 TeV protons from the circulating beam by means of a bent crystal.

... The idea is to put a bent, single crystal of either Si or Ge (W would perform slightly better but needs substantial improvements in crystal quality) at a distance of $\simeq 7\sigma$ to the beam where it can intercept and deflect part of the beam halo by an angle similar to the one the foreseen dump kicking system will apply to the circulating beam.

... ions with the same momentum per charge as protons are deflected in a crystal with similar efficiencies



If the crystal is positioned at the kicking section, the whole dump system can be used for slow extraction of parts of the beam halo, the particles that are anyway lost subsequently at collimators.


The beam extraction: news

[S. Montesano, *Physics at AFTER using LHC beams, ECT* Trento, Feb. 2013*]

Goal : assess the possibility to **use bent crystals as primary collimators** in hadronic accelerators and colliders



UA9 installation in the SPS




Prototype crystal collimation system at SPS :

- local **beam loss reduction** ($5\div 20\times$ reduction for proton beam)
- beam loss map show average loss reduction in the entire SPS ring
- **halo extraction efficiency**
 $70\div 80\%$ for protons ($50\div 70\%$ for Pb)

The beam extraction: news


[S. Montesano, *Physics at AFTER using LHC beams, ECT* Trento, Feb. 2013*]

Goal : assess the possibility to **use bent crystals as primary collimators** in hadronic accelerators and colliders



UA9 installation in the SPS

2010 - 2012



LUA9 future installation in LHC

Prototype crystal collimation system at SPS :

- local **beam loss reduction** ($5+20\times$ reduction for proton beam)
- beam loss map show average loss reduction in the entire SPS ring
- **halo extraction efficiency**
 $70+80\%$ for protons ($50+70\%$ for Pb)

The beam extraction: news


[S. Montesano, Physics at AFTER using LHC beams, ECT* Trento, Feb. 2013]

Goal : assess the possibility to **use bent crystals as primary collimators** in hadronic accelerators and colliders



UA9 installation in the SPS

2010 - 2012



LUA9 future installation in LHC

Prototype crystal collimation system at SPS :

- local **beam loss reduction** (5+20x reduction for proton beam)
- beam loss map show average loss reduction in the entire SPS ring
- **halo extraction efficiency**
70+80% for protons (50+70% for Pb)

Towards an installation in the LHC : propose and **install during LSI** a min. number of devices

- 2 crystals

Long term plan is ambitious : **propose a collimation system based on bent crystals** for the upgrade of the current LHC collimation system

Luminosities

- Instantaneous Luminosity:

$$\mathcal{L} = \Phi_{beam} \times N_{target} = N_{beam} \times (\rho \times \ell \times \mathcal{N}_A) / A$$

$$\Phi_{beam} = 2 \times 10^5 \text{ Pb s}^{-1}, \quad \ell = 1 \text{ cm (target thickness)}$$



- Integrated luminosity $\int dt \mathcal{L} = \mathcal{L} \times 10^6 \text{ s}$ for Pb
- Expected luminosities with $2 \times 10^5 \text{ Pb s}^{-1}$ extracted (1cm-long target)

Target	$\rho \text{ (g.cm}^{-3}\text{)}$	A	$\mathcal{L} \text{ (mb}^{-1}\text{.s}^{-1}\text{)} = \int \mathcal{L} \text{ (nb}^{-1}\text{.yr}^{-1}\text{)}$
Sol. H₂	0.09	1	11
Liq. H₂	0.07	1	8
Liq. D₂	0.16	2	10
Be	1.85	9	25
Cu	8.96	64	17
W	19.1	185	13
Pb	11.35	207	7

- Planned lumi for PHENIX Run15AuAu 2.8 nb^{-1} (0.13 nb^{-1} at 62 GeV)
- Nominal LHC lumi for PbPb 0.5 nb^{-1}

Crystal resistance to irradiation

- **IHEP U-70** (Biryukov et al, NIMB 234, 23-30):
 - 70 GeV protons, 50 ms spills of **10^{14} protons every 9.6 s**, several minutes irradiation
 - equivalent to 2 nominal LHC bunches for 500 turns every 10 s
 - 5 mm silicon crystal, **channeling efficiency unchanged**
- **SPS North Area - NA48** (Biino et al, CERN-SL-96-30-EA):
 - 450 GeV protons, 2.4 s spill of 5×10^{12} protons every 14.4 s, one year irradiation, **2.4×10^{20} protons/cm²** in total,
 - equivalent to several year of operation for a primary collimator in LHC
 - $10 \times 50 \times 0.9$ mm³ silicon crystal, 0.8×0.3 mm² area irradiated, **channeling efficiency reduced by 30%**.
- **HRMT16-UA9CRY** (HiRadMat facility, November 2012):
 - 440 GeV protons, up to 288 bunches in **7.2 μ s**, 1.1×10^{11} protons per bunch (**3×10^{13} protons** in total)
 - energy deposition comparable to an asynchronous beam dump in LHC
 - 3 mm long silicon crystal, **no damage to the crystal after accurate visual inspection**, more tests planned to assess possible crystal lattice damage
 - **accurate FLUKA simulation of energy deposition** and residual dose



AFTER, among other things, a quarkonium observatory in pp

- Interpolating the world data set:

Target	$\int \mathcal{L} \text{ (fb}^{-1}\text{.yr}^{-1}\text{)}$	$N(J/\Psi) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Upsilon}$
1 m Liq. H_2	20	$4.0 \cdot 10^8$	$8.0 \cdot 10^5$
1 m Liq. D_2	24	$9.6 \cdot 10^8$	$1.9 \cdot 10^6$
LHC pp 14 Tev (low pT)	0.05 (ALICE) 2 LHCb	$3.6 \cdot 10^7$ $1.4 \cdot 10^9$	$1.8 \cdot 10^5$ $7.2 \cdot 10^6$
RHIC pp 200GeV	$1.2 \cdot 10^{-2}$	$4.8 \cdot 10^5$	$1.2 \cdot 10^3$

AFTER, among other things, a quarkonium observatory in pp

- Interpolating the world data set:

Target	$\int \mathcal{L} \text{ (fb}^{-1}\text{.yr}^{-1}\text{)}$	$N(J/\Psi) \text{ yr}^{-1}$ $= A\mathcal{L}\mathcal{B}\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A\mathcal{L}\mathcal{B}\sigma_{\Upsilon}$
1 m Liq. H_2	20	$4.0 \cdot 10^8$	$8.0 \cdot 10^5$
1 m Liq. D_2	24	$9.6 \cdot 10^8$	$1.9 \cdot 10^6$
LHC pp 14 Tev (low pT)	0.05 (ALICE) 2 LHCb	$3.6 \cdot 10^7$ $1.4 \cdot 10^9$	$1.8 \cdot 10^5$ $7.2 \cdot 10^6$
RHIC pp 200GeV	$1.2 \cdot 10^{-2}$	$4.8 \cdot 10^5$	$1.2 \cdot 10^3$

- 1000 times higher than at RHIC; comparable to ALICE/LHCb at the LHC

AFTER, among other things, a quarkonium observatory in pp

- Interpolating the world data set:

Target	$\int \mathcal{L} \text{ (fb}^{-1}\cdot\text{yr}^{-1}\text{)}$	$N(J/\Psi) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Upsilon}$
1 m Liq. H_2	20	$4.0 \cdot 10^8$	$8.0 \cdot 10^5$
1 m Liq. D_2	24	$9.6 \cdot 10^8$	$1.9 \cdot 10^6$
LHC pp 14 Tev (low pT)	0.05 (ALICE) 2 LHCb	$3.6 \cdot 10^7$ $1.4 \cdot 10^9$	$1.8 \cdot 10^5$ $7.2 \cdot 10^6$
RHIC pp 200GeV	$1.2 \cdot 10^{-2}$	$4.8 \cdot 10^5$	$1.2 \cdot 10^3$

- 1000 times higher than at RHIC; comparable to ALICE/LHCb at the LHC
- Numbers are for only one unit of rapidity about 0

AFTER, among other things, a quarkonium observatory in pp

- Interpolating the world data set:

Target	$\int \mathcal{L} \text{ (fb}^{-1}\text{.yr}^{-1}\text{)}$	$N(J/\Psi) \text{ yr}^{-1}$ $= A \mathcal{L} B \sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A \mathcal{L} B \sigma_{\Upsilon}$
1 m Liq. H₂	20	4.0 10⁸	8.0 10⁵
1 m Liq. D₂	24	9.6 10⁸	1.9 10⁶
LHC pp 14 Tev (low pT)	0.05 (ALICE) 2 LHCb	3.6 10⁷ 1.4 10⁹	1.8 10⁵ 7.2 10⁶
RHIC pp 200GeV	1.2 10⁻²	4.8 10⁵	1.2 10³

- 1000 times higher than at RHIC; comparable to ALICE/LHCb at the LHC
- Numbers are for only one unit of rapidity about 0
- Unique access in the backward region

AFTER, among other things, a quarkonium observatory in pp

- Interpolating the world data set:

Target	$\int \mathcal{L} \text{ (fb}^{-1}\cdot\text{yr}^{-1})$	$N(J/\Psi) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Upsilon}$
1 m Liq. H_2	20	4.0 10^8	8.0 10^5
1 m Liq. D_2	24	9.6 10^8	1.9 10^6
LHC pp 14 Tev (low pT)	0.05 (ALICE) 2 LHCb	3.6 10^7 1.4 10^9	1.8 10^5 7.2 10^6
RHIC pp 200GeV	1.2 10^{-2}	4.8 10^5	1.2 10^3

- 1000 times higher than at RHIC; comparable to ALICE/LHCb at the LHC
- Numbers are for only one unit of rapidity about 0
- Unique access in the backward region
- Probe of the (very) large x in the target

Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**

Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**
 - in photo/lepto production (DIS)
 - but also pp collisions in gg -fusion process
 - mainly because of the presence of a natural “hard” scale: m_Q
 - and the good detectability of a dimuon pair

Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**
 - in photo/lepto production (DIS)
 - but also pp collisions in gg -fusion process
 - mainly because of the presence of a natural “hard” scale: m_Q
 - and the good detectability of a dimuon pair

PHYSICAL REVIEW D

VOLUME 37, NUMBER 5

1 MARCH 1988

Structure-function analysis and ψ , jet, W , and Z production: Determining the gluon distribution

A. D. Martin

Department of Physics, University of Durham, Durham, England

R. G. Roberts

Rutherford Appleton Laboratory, Didcot, Oxon, England

W. J. Stirling

Department of Physics, University of Durham, Durham, England

(Received 27 July 1987)

We perform a next-to-leading-order structure-function analysis of deep-inelastic μN and νN scattering data and find acceptable fits for a range of input gluon distributions. We show three equally acceptable sets of parton distributions which correspond to gluon distributions which are (1) “soft,” (2) “hard,” and (3) which behave as $xG(x) \sim 1/\sqrt{x}$ at small x . J/ψ and prompt photon hadroproduction data are used to discriminate between the three sets. Set 1, with the “soft”-gluon distribution, is favored. W , Z , and jet production data from the CERN collider are well described but do not distinguish between the sets of structure functions. The precision of the predictions for σ_W and σ_Z allow the collider measurements to yield information on the number of light neutrinos and the mass of the top quark. Finally we discuss how the gluon distribution at very small x may be directly measured at DESY HERA.

Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**
 - in photo/lepto production (DIS)
 - but also pp collisions in gg -fusion process
 - mainly because of the presence of a natural “hard” scale: m_Q
 - and the good detectability of a dimuon pair

PHYSICAL REVIEW D

VOLUME 37, NUMBER 5

1 MARCH 1988

Structure-function analysis and ψ , jet, W , and Z production: Determining the gluon distribution

A. D. Martin

Department of Physics, University of Durham, Durham, England

R. G. Roberts

Rutherford Appleton Laboratory, Didcot, Oxon, England

W. J. Stirling

Department of Physics, University of Durham, Durham, England

(Received 27 July 1987)

We perform a next-to-leading-order structure-function analysis of deep-inelastic μN and νN scattering data and find acceptable fits for a range of input gluon distributions. We show three equally acceptable sets of parton distributions which correspond to gluon distributions which are (1) “soft,” (2) “hard,” and (3) which behave as $xG(x) \sim 1/\sqrt{x}$ at small x . J/ψ and prompt photon hadroproduction data are used to discriminate between the three sets. Set 1, with the “soft”-gluon distribution, is favored. W , Z , and jet production data from the CERN collider are well described but do not distinguish between the sets of structure functions. The precision of the predictions for σ_W and σ_Z allow the collider measurements to yield information on the number of light neutrinos and the mass of the top quark. Finally we discuss how the gluon distribution at very small x may be directly measured at DESY HERA.

- Production **puzzle** → quarkonium not used anymore in global fits

Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**
 - in photo/lepto production (DIS)
 - but also pp collisions in gg -fusion process
 - mainly because of the presence of a natural “hard” scale: m_Q
 - and the good detectability of a dimuon pair

PHYSICAL REVIEW D

VOLUME 37, NUMBER 5

1 MARCH 1988

Structure-function analysis and ψ , jet, W , and Z production: Determining the gluon distribution

A. D. Martin

Department of Physics, University of Durham, Durham, England

R. G. Roberts

Rutherford Appleton Laboratory, Didcot, Oxon, England

W. J. Stirling

Department of Physics, University of Durham, Durham, England

(Received 27 July 1987)

We perform a next-to-leading-order structure-function analysis of deep-inelastic μN and νN scattering data and find acceptable fits for a range of input gluon distributions. We show three equally acceptable sets of parton distributions which correspond to gluon distributions which are (1) “soft,” (2) “hard,” and (3) which behave as $xG(x) \sim 1/\sqrt{x}$ at small x . J/ψ and prompt photon hadroproduction data are used to discriminate between the three sets. Set 1, with the “soft”-gluon distribution, is favored. W , Z , and jet production data from the CERN collider are well described but do not distinguish between the sets of structure functions. The precision of the predictions for σ_W and σ_Z allow the collider measurements to yield information on the number of light neutrinos and the mass of the top quark. Finally we discuss how the gluon distribution at very small x may be directly measured at DESY HERA.

- Production **puzzle** \rightarrow quarkonium not used anymore in global fits
- With systematic studies, one would **restore its status as gluon probe**

Accessing the large x glue with quarkonia

PYTHIA simulation
 $\sigma(y) / \sigma(y=0.4)$
 statistics for one month
 5% acceptance considered

Statistical relative uncertainty
 Large statistics allow to access
 very backward region

Gluon uncertainty from
 MSTWPDF
 - only for the gluon content of
 the target
 - assuming

$$x_g = M_{J/\psi} / \sqrt{s} e^{-y_{CM}}$$

J/ψ


$$y_{CM} \sim 0 \rightarrow x_g = 0.03$$

$$y_{CM} \sim -3.6 \rightarrow x_g = 1$$

Y : larger x_g for same y_{CM}

$$y_{CM} \sim 0 \rightarrow x_g = 0.08$$

$$y_{CM} \sim -2.4 \rightarrow x_g = 1$$



\Rightarrow Backward measurements allow to access large x gluon pdf


NEW!

(x, Q^2) map of AFTER isolated- γ


[D.d'E & J.Rojo, NPB 860 (2012) 311]

■ p-p kinematics at fixed-target LHC:

To access $x > 0.3$ one needs isolated- γ with: $p_T = x_T \sqrt{s}/2 > 10\text{-}20 \text{ GeV}/c$



AFTER region: $pp \rightarrow \gamma X$



[D. D'Enterria, Physics at AFTER using LHC beams, ECT* Trento, Feb 2013]

AFTER: also a quarkonium observatory in pA

Target	A	$\int \mathcal{L} \text{ (fb}^{-1}\text{.yr}^{-1}\text{)}$	$N(J/\Psi) \text{ yr}^{-1}$ $= A\mathcal{L}\mathcal{B}\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A\mathcal{L}\mathcal{B}\sigma_{\Upsilon}$
1cm Be	9	0.62	$1.1 \cdot 10^8$	$2.2 \cdot 10^5$
1cm Cu	64	0.42	$5.3 \cdot 10^8$	$1.1 \cdot 10^6$
1cm W	185	0.31	$1.1 \cdot 10^9$	$2.3 \cdot 10^6$
1cm Pb	207	0.16	$6.7 \cdot 10^8$	$1.3 \cdot 10^6$
LHC pPb 8.8 TeV	207	10^{-4}	$1.0 \cdot 10^7$	$7.5 \cdot 10^4$
RHIC dAu 200GeV	198	$1.5 \cdot 10^{-4}$	$2.4 \cdot 10^6$	$5.9 \cdot 10^3$
RHIC dAu 62GeV	198	$3.8 \cdot 10^{-6}$	$1.2 \cdot 10^4$	18

- In principle, one can get **300 times more J/ψ** —not counting the likely wider y coverage— than at RHIC, allowing for

AFTER: also a quarkonium observatory in pA

Target	A	$\int \mathcal{L} \text{ (fb}^{-1}\text{.yr}^{-1}\text{)}$	$N(J/\Psi) \text{ yr}^{-1}$ $= A \mathcal{L} B \sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A \mathcal{L} B \sigma_{\Upsilon}$
1cm Be	9	0.62	$1.1 \cdot 10^8$	$2.2 \cdot 10^5$
1cm Cu	64	0.42	$5.3 \cdot 10^8$	$1.1 \cdot 10^6$
1cm W	185	0.31	$1.1 \cdot 10^9$	$2.3 \cdot 10^6$
1cm Pb	207	0.16	$6.7 \cdot 10^8$	$1.3 \cdot 10^6$
LHC pPb 8.8 TeV	207	10^{-4}	$1.0 \cdot 10^7$	$7.5 \cdot 10^4$
RHIC dAu 200GeV	198	$1.5 \cdot 10^{-4}$	$2.4 \cdot 10^6$	$5.9 \cdot 10^3$
RHIC dAu 62GeV	198	$3.8 \cdot 10^{-6}$	$1.2 \cdot 10^4$	18

- In principle, one can get **300 times more J/ψ** –not counting the likely wider y coverage– than at RHIC, allowing for
 - χ_c measurement in pA via $J/\psi + \gamma$ (extending Hera-B studies)

AFTER: also a quarkonium observatory in pA

Target	A	$\int \mathcal{L} \text{ (fb}^{-1}\cdot\text{yr}^{-1})$	$N(J/\Psi) \text{ yr}^{-1}$ $= A\mathcal{L}\mathcal{B}\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A\mathcal{L}\mathcal{B}\sigma_{\Upsilon}$
1cm Be	9	0.62	$1.1 \cdot 10^8$	$2.2 \cdot 10^5$
1cm Cu	64	0.42	$5.3 \cdot 10^8$	$1.1 \cdot 10^6$
1cm W	185	0.31	$1.1 \cdot 10^9$	$2.3 \cdot 10^6$
1cm Pb	207	0.16	$6.7 \cdot 10^8$	$1.3 \cdot 10^6$
LHC pPb 8.8 TeV	207	10^{-4}	$1.0 \cdot 10^7$	$7.5 \cdot 10^4$
RHIC dAu 200GeV	198	$1.5 \cdot 10^{-4}$	$2.4 \cdot 10^6$	$5.9 \cdot 10^3$
RHIC dAu 62GeV	198	$3.8 \cdot 10^{-6}$	$1.2 \cdot 10^4$	18

- In principle, one can get **300 times more J/ψ** —not counting the likely wider y coverage— than at RHIC, allowing for
 - χ_c measurement in pA via $J/\psi + \gamma$ (extending Hera-B studies)
 - **Polarisation** measurement as **the centrality, y or P_T**

AFTER: also a quarkonium observatory in pA

Target	A	$\int \mathcal{L} \text{ (fb}^{-1}\cdot\text{yr}^{-1})$	$N(J/\Psi) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Upsilon}$
1cm Be	9	0.62	$1.1 \cdot 10^8$	$2.2 \cdot 10^5$
1cm Cu	64	0.42	$5.3 \cdot 10^8$	$1.1 \cdot 10^6$
1cm W	185	0.31	$1.1 \cdot 10^9$	$2.3 \cdot 10^6$
1cm Pb	207	0.16	$6.7 \cdot 10^8$	$1.3 \cdot 10^6$
LHC pPb 8.8 TeV	207	10^{-4}	$1.0 \cdot 10^7$	$7.5 \cdot 10^4$
RHIC dAu 200GeV	198	$1.5 \cdot 10^{-4}$	$2.4 \cdot 10^6$	$5.9 \cdot 10^3$
RHIC dAu 62GeV	198	$3.8 \cdot 10^{-6}$	$1.2 \cdot 10^4$	18

- In principle, one can get **300 times more J/ψ** –not counting the likely wider y coverage– than at RHIC, allowing for
 - χ_c measurement in pA via $J/\psi + \gamma$ (extending Hera-B studies)
 - **Polarisation** measurement as **the centrality, y or P_T**
 - Ratio ψ' over **direct J/ψ** measurement in pA

AFTER: also a quarkonium observatory in pA

Target	A	$\int \mathcal{L} \text{ (fb}^{-1}\cdot\text{yr}^{-1})$	$N(J/\Psi) \text{ yr}^{-1}$ $= A\mathcal{L}\mathcal{B}\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A\mathcal{L}\mathcal{B}\sigma_{\Upsilon}$
1cm Be	9	0.62	$1.1 \cdot 10^8$	$2.2 \cdot 10^5$
1cm Cu	64	0.42	$5.3 \cdot 10^8$	$1.1 \cdot 10^6$
1cm W	185	0.31	$1.1 \cdot 10^9$	$2.3 \cdot 10^6$
1cm Pb	207	0.16	$6.7 \cdot 10^8$	$1.3 \cdot 10^6$
LHC pPb 8.8 TeV	207	10^{-4}	$1.0 \cdot 10^7$	$7.5 \cdot 10^4$
RHIC dAu 200GeV	198	$1.5 \cdot 10^{-4}$	$2.4 \cdot 10^6$	$5.9 \cdot 10^3$
RHIC dAu 62GeV	198	$3.8 \cdot 10^{-6}$	$1.2 \cdot 10^4$	18

- In principle, one can get **300 times more J/ψ** —not counting the likely wider y coverage— than at RHIC, allowing for
 - χ_c measurement in pA via $J/\psi + \gamma$ (extending Hera-B studies)
 - **Polarisation** measurement as **the centrality, y or P_T**
 - Ratio ψ' over **direct J/ψ** measurement in pA
 - not to mention ratio with **open charm, Drell-Yan**, etc ...

What for ?

- The **target versatility** of a fixed-target experiment is undisputable

What for ?

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
 - a precise analysis of **gluon nuclear PDF**: $y, p_T \leftrightarrow x_2$
 - a handle on **formation time effects**

What for ?

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
 - a precise analysis of **gluon nuclear PDF**: $y, p_T \leftrightarrow x_2$
 - a handle on **formation time effects**
- Strong need for **cross checks** from **various** measurements

What for ?

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
 - a precise analysis of **gluon nuclear PDF**: $y, p_T \leftrightarrow x_2$
 - a handle on **formation time effects**
- Strong need for **cross checks** from **various** measurements
- The **backward kinematics** is very useful for large- x_{target} studies

What for ?

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
 - a precise analysis of **gluon nuclear PDF**: $y, p_T \leftrightarrow x_2$
 - a handle on **formation time effects**
- Strong need for **cross checks** from **various** measurements
- The **backward kinematics** is very useful for large- x_{target} studies
 - What is the amount of Intrinsic charm ? Is it color filtered ?

What for ?

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
 - a precise analysis of **gluon nuclear PDF**: $y, p_T \leftrightarrow x_2$
 - a handle on **formation time effects**
- Strong need for **cross checks** from **various** measurements
- The **backward kinematics** is very useful for large- x_{target} studies
 - What is the amount of Intrinsic charm ? Is it color filtered ?
 - **Is there an EMC effect for gluon ?** (reminder: EMC region $0.3 < x < 0.7$)

What for ?

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
 - a precise analysis of **gluon nuclear PDF**: $y, p_T \leftrightarrow x_2$
 - a handle on **formation time effects**
- Strong need for **cross checks** from **various** measurements
- The **backward kinematics** is very useful for large- x_{target} studies
 - What is the amount of Intrinsic charm ? Is it color filtered ?
 - **Is there an EMC effect for gluon ?** (reminder: EMC region $0.3 < x < 0.7$)
- One should be careful with factorization breaking effects:

This calls for **multiple measurements** to (in)validate factorization

AFTER: also an heavy-flavour observatory in PbA

- Luminosities and yields with the extracted 2.76 TeV Pb beam
($\sqrt{s_{NN}} = 72$ GeV)

Target	A.B	$\int \mathcal{L} \text{ (nb}^{-1}\text{.yr}^{-1}\text{)}$	$N(J/\Psi) \text{ yr}^{-1}$ $= AB \mathcal{L} B \sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= AB \mathcal{L} B \sigma_{\Upsilon}$
1 m Liq. H₂	207.1	800	3.4 10⁶	6.9 10³
1cm Be	207.9	25	9.1 10⁵	1.9 10³
1cm Cu	207.64	17	4.3 10⁶	0.9 10³
1cm W	207.185	13	9.7 10⁶	1.9 10⁴
1cm Pb	207.207	7	5.7 10⁶	1.1 10⁴
LHC PbPb 5.5 TeV	207.207	0.5	7.3 10⁶	3.6 10⁴
RHIC AuAu 200GeV	198.198	2.8	4.4 10⁶	1.1 10⁴
RHIC AuAu 62GeV	198.198	0.13	4.0 10⁴	61

AFTER: also an heavy-flavour observatory in PbA

- Luminosities and yields with the extracted 2.76 TeV Pb beam
($\sqrt{s_{NN}} = 72$ GeV)

Target	A.B	$\int \mathcal{L} \text{ (nb}^{-1}\text{.yr}^{-1}\text{)}$	$N(J/\Psi) \text{ yr}^{-1}$ $= AB\mathcal{L}B\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= AB\mathcal{L}B\sigma_{\Upsilon}$
1 m Liq. H₂	207.1	800	3.4 10⁶	6.9 10³
1cm Be	207.9	25	9.1 10⁵	1.9 10³
1cm Cu	207.64	17	4.3 10⁶	0.9 10³
1cm W	207.185	13	9.7 10⁶	1.9 10⁴
1cm Pb	207.207	7	5.7 10⁶	1.1 10⁴
LHC PbPb 5.5 TeV	207.207	0.5	7.3 10⁶	3.6 10⁴
RHIC AuAu 200GeV	198.198	2.8	4.4 10⁶	1.1 10⁴
RHIC AuAu 62GeV	198.198	0.13	4.0 10⁴	61

- Yields **similar** to those of RHIC at 200 GeV,
100 times those of RHIC at 62 GeV

AFTER: also an heavy-flavour observatory in PbA

- Luminosities and yields with the extracted 2.76 TeV Pb beam
($\sqrt{s_{NN}} = 72$ GeV)

Target	A.B	$\int \mathcal{L} \text{ (nb}^{-1}\text{.yr}^{-1}\text{)}$	$N(J/\Psi) \text{ yr}^{-1}$ $= AB\mathcal{L}B\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= AB\mathcal{L}B\sigma_{\Upsilon}$
1 m Liq. H₂	207.1	800	3.4 10⁶	6.9 10³
1cm Be	207.9	25	9.1 10⁵	1.9 10³
1cm Cu	207.64	17	4.3 10⁶	0.9 10³
1cm W	207.185	13	9.7 10⁶	1.9 10⁴
1cm Pb	207.207	7	5.7 10⁶	1.1 10⁴
LHC PbPb 5.5 TeV	207.207	0.5	7.3 10⁶	3.6 10⁴
RHIC AuAu 200GeV	198.198	2.8	4.4 10⁶	1.1 10⁴
RHIC AuAu 62GeV	198.198	0.13	4.0 10⁴	61

- Yields **similar** to those of RHIC at 200 GeV,
100 times those of RHIC at 62 GeV
- Also **very competitive** compared to the **LHC**.

AFTER: also an heavy-flavour observatory in PbA

- Luminosities and yields with the extracted 2.76 TeV Pb beam
($\sqrt{s_{NN}} = 72$ GeV)

Target	A.B	$\int \mathcal{L} \text{ (nb}^{-1}\text{.yr}^{-1}\text{)}$	$N(J/\Psi) \text{ yr}^{-1}$ $= AB\mathcal{L}B\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= AB\mathcal{L}B\sigma_{\Upsilon}$
1 m Liq. H₂	207.1	800	3.4 10⁶	6.9 10³
1cm Be	207.9	25	9.1 10⁵	1.9 10³
1cm Cu	207.64	17	4.3 10⁶	0.9 10³
1cm W	207.185	13	9.7 10⁶	1.9 10⁴
1cm Pb	207.207	7	5.7 10⁶	1.1 10⁴
LHC PbPb 5.5 TeV	207.207	0.5	7.3 10⁶	3.6 10⁴
RHIC AuAu 200GeV	198.198	2.8	4.4 10⁶	1.1 10⁴
RHIC AuAu 62GeV	198.198	0.13	4.0 10⁴	61

- Yields **similar** to those of RHIC at 200 GeV,
100 times those of RHIC at 62 GeV
- Also **very competitive** compared to the **LHC**.

The same picture also holds for **open heavy flavour**

What for ?

Observation of J/ψ sequential suppression **seems to be hindered** by

- the **Cold Nuclear Matter effects**: non trivial and
... not well understood

What for ?


- Observation of J/ψ sequential suppression **seems to be hindered** by
- the **Cold Nuclear Matter effects**: non trivial and
... not well understood
 - the difficulty to observe directly the **excited states**
which would melt before the ground states
 - χ_c **never studied in AA** collisions
 - $\psi(2S)$ **not yet** studied in AA collisions **at RHIC**

What for ?

- Observation of J/ψ sequential suppression **seems to be hindered** by
- the **Cold Nuclear Matter effects**: non trivial and
... not well understood
 - the difficulty to observe directly the **excited states**
which would melt before the ground states
 - χ_c **never studied in AA** collisions
 - $\psi(2S)$ **not yet** studied in AA collisions **at RHIC**
 - the possibilities for **$c\bar{c}$ recombination**
 - Open charm** studies are **difficult** where recombination matters most
i.e. at **low P_T**
 - Only indirect indications –from the y and P_T dependence of R_{AA} –
that recombination may be at work
 - CNM effects may show a non-trivial y and P_T dependence ...

SPS and Hera-B


– J/ψ data in pA collisions



NA60 Phys.Lett. B 706 (2012) 263
 NA 50 Eur.Phys.J. C48 (2006) 329
 NA 3 Z.Phys. C20 (1983)
 HERA-B Eur.Phys.J. C60 (2009) 525


SPS and Hera-B

– J/ψ data in pA collisions



NA60 Phys.Lett. B 706 (2012) 263
 NA 50 Eur.Phys.J. C48 (2006) 329
 NA 3 Z.Phys. C20 (1983)
 HERA-B Eur.Phys.J. C60 (2009) 525

– χ_c data in pA collisions




HERA-B PRD 79 (2009) 012001, and ref. therein

A Fixed Target Experiment

A dilepton observatory

→ Region in x probed by dilepton production as function of $M_{\ell\ell}$




A Fixed Target Experiment

A dilepton observatory

→ Region in x probed by dilepton production as function of $M_{\ell\ell}$

→ Above $c\bar{c}$: $x \in [10^{-3}, 1]$

→ Above $b\bar{b}$: $x \in [9 \times 10^{-3}, 1]$




A Fixed Target Experiment

A dilepton observatory

→ Region in x probed by dilepton production as function of $M_{\ell\ell}$

→ Above $c\bar{c}$: $x \in [10^{-3}, 1]$

→ Above $b\bar{b}$: $x \in [9 \times 10^{-3}, 1]$



A Fixed Target Experiment


A dilepton observatory

→ Region in x probed by dilepton production as function of $M_{\ell\ell}$

→ Above $c\bar{c}$: $x \in [10^{-3}, 1]$

→ Above $b\bar{b}$: $x \in [9 \times 10^{-3}, 1]$

Note: $x_{\text{target}}(\equiv x_2) > x_{\text{projectile}}(\equiv x_1)$
“backward” region



A Fixed Target Experiment

A dilepton observatory

→ Region in x probed by dilepton production as function of $M_{\ell\ell}$


→ Above $c\bar{c}$: $x \in [10^{-3}, 1]$

→ Above $b\bar{b}$: $x \in [9 \times 10^{-3}, 1]$

Note: $x_{\text{target}} (\equiv x_2) > x_{\text{projectile}} (\equiv x_1)$
“backward” region

→ sea-quark asymmetries
via p and d studies

- at large(est) x : backward (“easy”)
- at small(est) x : forward (need to stop the (extracted) beam)



A Fixed Target Experiment

A dilepton observatory

⇒ Region in x probed by dilepton production as function of $M_{\ell\ell}$


→ Above $c\bar{c}$: $x \in [10^{-3}, 1]$

→ Above $b\bar{b}$: $x \in [9 \times 10^{-3}, 1]$

Note: $x_{\text{target}} (\equiv x_2) > x_{\text{projectile}} (\equiv x_1)$
“backward” region

→ **sea-quark asymmetries**
via p and d studies

- at large(est) x : backward (“easy”)
- at small(est) x : forward (need to stop the (extracted) beam)



⇒ To do: to look at the rates to see how competitive this will be

A Fixed Target Experiment

SSA in Drell-Yan studies

⇒ Relevant parameters for the future **planned polarized DY experiments**.

Experiment	particles	energy (GeV)	\sqrt{s} (GeV)	x_p^\uparrow	\mathcal{L} (nb ⁻¹ s ⁻¹)
AFTER	$p + p^\uparrow$	7000	115	$0.01 \div 0.9$	1
COMPASS	$\pi^\pm + p^\uparrow$	160	17.4	$0.2 \div 0.3$	2
COMPASS (low mass)	$\pi^\pm + p^\uparrow$	160	17.4	~ 0.05	2
RHIC	$p^\uparrow + p$	collider	500	$0.05 \div 0.1$	0.2
J-PARC	$p^\uparrow + p$	50	10	$0.5 \div 0.9$	1000
PANDA (low mass)	$\bar{p} + p^\uparrow$	15	5.5	$0.2 \div 0.4$	0.2
PAX	$p^\uparrow + \bar{p}$	collider	14	$0.1 \div 0.9$	0.002
NICA	$p^\uparrow + p$	collider	20	$0.1 \div 0.8$	0.001
RHIC	$p^\uparrow + p$	250	22	$0.2 \div 0.5$	2
Int.Target 1					
RHIC	$p^\uparrow + p$	250	22	$0.2 \div 0.5$	60
Int.Target 2					

⇒ For AFTER, numbers correspond to a 50 cm polarized H target.

⇒ $\ell^+\ell^-$ angular distribution: separation Siverts vs. Boer-Mulders effects

LHB

Our idea is not completely new

Nuclear Instruments and Methods in Physics Research A 333 (1993) 125–135
North-Holland

**NUCLEAR
INSTRUMENTS
& METHODS
IN PHYSICS
RESEARCH**
Section A

LHB, a fixed target experiment at LHC to measure CP violation in B mesons

Flavio Costantini

University of Pisa and INFN, Italy

A fixed target experiment at LHC to measure CP violation in B mesons is presented. A description of the proposed apparatus is given together with its sensitivity on the CP violation asymmetry measurement for the two benchmark decay channels $B^0 \rightarrow J/\psi + K_s^0$, $B^0 \rightarrow \pi^+ \pi^-$. The possibility of obtaining an extracted LHC beam hinges on channeling in a bent silicon crystal. Recent results on beam extraction efficiencies measured at CERN SPS based on this technique are presented.

LHB

Our idea is not completely new

1. Introduction

...

This paper presents a fixed target experiment to measure CP violation in the B system based on the possibility of extracting the 8 TeV LHC proton beam using a bent silicon crystal [4]. A 10% extraction efficiency of the LHC beam halo will give an extracted beam intensity of about 10^8 protons/s allowing the production of as many as 10^{10} $B\bar{B}$ pairs per year, i.e. about two orders of magnitude more than what could be produced by an e^+e^- asymmetric B factory with 10^{34} $\text{cm}^{-2}\text{s}^{-1}$ luminosity [5].



LHB

Our idea is not completely new

1. Introduction

...

This paper presents a fixed target experiment to measure CP violation in the B system based on the possibility of extracting the 8 TeV LHC proton beam using a bent silicon crystal [4]. A 10% extraction efficiency of the LHC beam halo will give an extracted beam intensity of about 10^8 protons/s allowing the production of as many as 10^{10} $B\bar{B}$ pairs per year, i.e. about two orders of magnitude more than what could be produced by an e^+e^- asymmetric B factory with 10^{34} $\text{cm}^{-2}\text{s}^{-1}$ luminosity [5].

10^{10} $B\bar{B}$ pairs per year



- B -factories: 1 ab^{-1} means $10^9 B\bar{B}$ pairs

LHB

Our idea is not completely new

1. Introduction

...

This paper presents a fixed target experiment to measure CP violation in the B system based on the possibility of extracting the 8 TeV LHC proton beam using a bent silicon crystal [4]. A 10% extraction efficiency of the LHC beam halo will give an extracted beam intensity of about 10^8 protons/s allowing the production of as many as 10^{10} $B\bar{B}$ pairs per year, i.e. about two orders of magnitude more than what could be produced by an e^+e^- asymmetric B factory with $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ luminosity [5].

10^{10} $B\bar{B}$ pairs per year



- B-factories: 1 ab^{-1} means $10^9 B\bar{B}$ pairs
- For LHCb, typically 1 fb^{-1} means $\simeq 2 \times 10^{11} B\bar{B}$ pairs at 14 TeV

LHB

Our idea is not completely new

1. Introduction

...

This paper presents a fixed target experiment to measure CP violation in the B system based on the possibility of extracting the 8 TeV LHC proton beam using a bent silicon crystal [4]. A 10% extraction efficiency of the LHC beam halo will give an extracted beam intensity of about 10^8 protons/s allowing the production of as many as 10^{10} $B\bar{B}$ pairs per year, i.e. about two orders of magnitude more than what could be produced by an e^+e^- asymmetric B factory with 10^{34} $\text{cm}^{-2}\text{s}^{-1}$ luminosity [5].

10^{10} $B\bar{B}$ pairs per year



- B-factories: 1 ab^{-1} means $10^9 B\bar{B}$ pairs
- For LHCb, typically 1 fb^{-1} means $\simeq 2 \times 10^{11} B\bar{B}$ pairs at 14 TeV
- LHB turned down in favour of LHCb mainly because of the **fear of a premature degradation of the bent crystal** due to radiation damages.

LHB

Our idea is not completely new

1. Introduction

...

This paper presents a fixed target experiment to measure CP violation in the B system based on the possibility of extracting the 8 TeV LHC proton beam using a bent silicon crystal [4]. A 10% extraction efficiency of the LHC beam halo will give an extracted beam intensity of about 10^8 protons/s allowing the production of as many as 10^{10} $B\bar{B}$ pairs per year, i.e. about two orders of magnitude more than what could be produced by an e^+e^- asymmetric B factory with $10^{34} \text{ cm}^{-2}\text{s}^{-1}$ luminosity [5].

10^{10} $B\bar{B}$ pairs per year



- B-factories: 1 ab^{-1} means $10^9 B\bar{B}$ pairs
- For LHCb, typically 1 fb^{-1} means $\simeq 2 \times 10^{11} B\bar{B}$ pairs at 14 TeV
- LHB turned down in favour of LHCb mainly because of the **fear of a premature degradation of the bent crystal** due to radiation damages.
- Nowadays, degradation is known to be $\simeq 6\%$ per 10^{20} particles/cm²
- 10^{20} particles/cm² : one year of operation for realistic conditions

LHB

Our idea is not completely new

1. Introduction

...

This paper presents a fixed target experiment to measure CP violation in the B system based on the possibility of extracting the 8 TeV LHC proton beam using a bent silicon crystal [4]. A 10% extraction efficiency of the LHC beam halo will give an extracted beam intensity of about 10^8 protons/s allowing the production of as many as 10^{10} $B\bar{B}$ pairs per year, i.e. about two orders of magnitude more than what could be produced by an e^+e^- asymmetric B factory with 10^{34} $\text{cm}^{-2}\text{s}^{-1}$ luminosity [5].

10^{10} $B\bar{B}$ pairs per year



- B-factories: 1 ab^{-1} means $10^9 B\bar{B}$ pairs
- For LHCb, typically 1 fb^{-1} means $\simeq 2 \times 10^{11} B\bar{B}$ pairs at 14 TeV
- LHB turned down in favour of LHCb mainly because of the **fear of a premature degradation of the bent crystal** due to radiation damages.
- Nowadays, degradation is known to be $\simeq 6\%$ per 10^{20} particles/ cm^2
- 10^{20} particles/ cm^2 : one year of operation for realistic conditions
- After a year, one simply moves the crystal by less than one mm ...