


AFTER@LHC: A Fixed Target Experiment *for hadron, heavy-ion and spin physics:* *Status and short-range plan*

Barbara Trzeciak

*Faculty of Nuclear Sciences and Physical Engineering
Czech Technical University in Prague*

M. Anselmino Dip. di Fisica and INFN Sez. Torino, Via P. Giuria 1, I-10125, Torino, Italy
R. Arnaldi and **E. Scomparin** INFN Sez. Torino, Via P. Giuria 1, I-10125, Torino, Italy
S.J. Brodsky and **C. Lorcé** SLAC National Accelerator Laboratory, Theoretical Physics, Stanford University, Menlo Park, CA 94025, USA
V. Chambert, **J.P. Didelez**, **B. Genolini**, **C. Hadjidakis**, **I. Hrivnacova**, and **J.P. Lansberg** IPNO, Université Paris-Sud, CNRS/IN2P3, F-91406, Orsay, France
E.G. Ferreira Departamento de Física de Partículas, Universidade de Santiago de Compostela, 15782 Santiago de Compostela, Spain
F. Fleuret Laboratoire Leprince Ringuet, Ecole Polytechnique, CNRS/IN2P3, 91128 Palaiseau, France
Y. Gao and **Z. Yang** CHEP, Department of Engineering Physics, Tsinghua University, Beijing, China
L. Massacrier LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France and IPNO, Université Paris-Sud, CNRS/IN2P3, F-91406, Orsay, France
R. Mikkelsen and **U.I. Uggerhøj** Department of Physics and Astronomy, University of Aarhus, Denmark
C. Pisano Antwerp U., Belgium
I. Schienbein LPSC, Université Joseph Fourier, CNRS/IN2P3/INPG, F-380 26 Grenoble, France
M. Schlegel Institute for Theoretical Physics, Tübingen U., D-72076 Tübingen, German
B. Trzeciak FNSPE, Czech Technical University, Prague, Czech Republic
R. Ulrich Institut für Kernphysik, Karlsruhe Institute of Technology (KIT), 76021 Karlsruhe, Germany



INVESTICE DO ROZVOJE VZDĚLÁVÁNÍ

EPS HEP
22-29 July, 2015
Vienna, Austria



Outline

- ✓ Advantages of a fixed target experiment at LHC
- ✓ Internal gas target vs beam extraction with a bent crystal
- ✓ Expected luminosities
- ✓ Physics Highlights
- ✓ Feasibility studies of quarkonium production



Outline

- ✓ Advantages of a fixed target experiment at LHC
- ✓ Internal gas target vs beam extraction with a bent crystal
- ✓ Expected luminosities
- ✓ Physics Highlights
- ✓ Feasibility studies of quarkonium production

WHAT IS AFTER@LHC AND WHAT FOR?

AFTER@LHC is a proposal for a multi-purpose fixed target experiment using the multi-TeV proton or heavy ion beams of the LHC

- **Advance our understanding of the large-x gluon antiquark and heavy-quark content in the nucleon and nucleus**
- **Dynamics and spin of gluons inside (un)polarised nucleons**
- **Heavy-ion collisions towards large rapidities**



Advantages of a fixed target experiment at LHC

✓ Advantages of a fixed-target experiment:

- high luminosities with dense targets
- target versatility
- possibility to polarize target
 - spin physics program
- access to large Feynman $|x_F|$

→ With LHC beams:

7 TeV proton beam on a fixed target

CMS energy: $\sqrt{s} = \sqrt{2m_N E_p} \approx 115 \text{ GeV}$	Rapidity shift: $y_{CM} = 0 \rightarrow y_{lab} = 4.8$
Boost: $\gamma = \sqrt{s} / (2m_p) \approx 60$	


2.76 TeV Pb beam on a fixed target

CMS energy: $\sqrt{s_{NN}} = \sqrt{2m_N E_{Pb}} \approx 72 \text{ GeV}$	Rapidity shift: $y_{CM} = 0 \rightarrow y_{lab} = 4.3$
Boost: $\gamma \approx 40$	



Advantages of a fixed target experiment at LHC

- ✓ Testing QCD at large $x = (0.3, 1)$




- ✓ Entire forward hemisphere – $y_{CM} > 0$ – within: $0^\circ < \theta_{lab} < 1^\circ$ - large occupancy – more challenging
- ✓ Backward region - $y_{CM} < 0$ – at large angles in the lab frame – low occupancy, no constrain from a beam pipe
 - Backward physics accessible
 - Access to partons with momentum fraction $x_2 \rightarrow 1$ in the target

($x_F \rightarrow -1$)



Beam extraction using bent crystal

✓ Possible fixed-target mode



- ✓ UA9: test @SPS on the crystal with proton and ion beams LUA9 (beam bending experiment using crystal): approved by LHCC
 - *2 bent crystals installed in IR7 during LS1, 2015/2016 first tests with beams*

➤ Proton beam extraction:

- Single or multi-pass extraction efficiency of 50%
- LHC beam loss $\sim 10^9 p^+ s^{-1}$ - extracted beam : $5 \times 10^8 p^+ s^{-1}$

➤ Ion beam extraction


- Successfully tested at the SPS, should also work at the LHC (P. Ballin et al, NIMB 267 (2009) 2952)



Beam extraction using bent crystal



✓ Possible fixed-target mode



- ✓ UA9: test @SPS on the crystal with proton and ion beams LUA9 (beam bending experiment using crystal): approved by LHCC
 - *2 bent crystals installed in IR7 during LS1, 2015/2016 first tests with beams*

➤ Proton beam extraction:

- Single or multi-pass extraction efficiency of 50%
- LHC beam loss $\sim 10^9 p^+ s^{-1}$ - extracted beam : $5 \times 10^8 p^+ s^{-1}$

➤ Ion beam extraction

- Successfully tested at the SPS, should also work at the LHC (P. Ballin et al, NIMB 267 (2009) 2952)

➤ Deflecting the beam halo at 7σ distance to the beam


➔ No loss in the LHC beam



Internal gas target, *SMOG@LHC*

✓ Possible fixed-target mode

SMOG: System for Measuring Overlap with Gas



- ✓ Low density Ne-gas injected into VELO in LHCb
- ✓ Short pNe pilot run at $\sqrt{s_{NN}} = 87$ GeV in 2012

LHCb-CONF-2012-034

- ✓ Short PbNe pilot run at $\sqrt{s_{NN}} = 54$ GeV in 2013

Ne target density: $1.5 \cdot 10^{-7}$ mbar

→ injection of Ne-gas into VELO

- Noble gases favored
- As for now, target polarization is not possible with SMOG
- Internal gas target can be polarized, would be another system with respect to SMOG



Luminosities in pH and pA at $\sqrt{s}_{NN} = 115 \text{ GeV}$

With bent crystal

- ✓ Instantaneous luminosity: $L = \phi_{\text{beam}} \times N_{\text{target}} = \phi_{\text{beam}} \times (\rho \times l \times N_A) / A$
 l is a target thickness
- ✓ $\phi_{\text{beam}} = 5 \times 10^8 \text{ p}^+ \text{ s}^{-1}$ (50% of the beam loss)
- ✓ Integrated luminosity - LHC year – 9 months running = 10^7 s

Target	ρ (g.cm ⁻³)	A	L ($\mu\text{b}^{-1}\text{s}^{-1}$)	$\int L$ (pb ⁻¹ yr ⁻¹)
Liq H ₂ (1m)	0.07	1	2000	20000
Liq D ₂ (1m)	0.16	2	2400	24000
Be (1cm)	1.85	9	62	620
Cu (1cm)	8.96	64	42	420
W (1cm)	19.1	185	31	310
Pb (1cm)	11.35	207	16	160

- Large luminosities comparable to LHC - with 1 m long H₂(D₂) target,
3 orders of magnitude larger than at RHIC



Luminosities in pA - bent crystal vs SMOG

With bent crystal

SMOG
based on the pilot run

Target	ρ (g.cm ⁻³)	A	L ($\mu\text{b}^{-1}\text{s}^{-1}$)	\int L (pb ⁻¹ yr ⁻¹)
Be (1cm)	1.85	9	62	620
Cu (1cm)	8.96	64	42	420

Target: Ne gas

- Ne target density: 10^{-6} mbar
- $L = 8 \mu\text{b}^{-1}\text{s}^{-1}$

- ✓ Higher **instantaneous** luminosities using a bent crystal compare to what is expected from SMOG from the pilot run - $62 \mu\text{b}^{-1}\text{s}^{-1}$ with 1cm Be target vs $8 \mu\text{b}^{-1}\text{s}^{-1}$ for Ne in SMOG
- ✓ Higher Ne pressure needed in SMOG in order to reach comparable luminosity as in the bent crystal case
 - ✓ assuming *1 year of running with a proton beam and $P \approx 10^{-5}$ mbar, one can obtain comparable luminosity as in the bent crystal case*
- Increasing the pressure is not expected to decrease the beam life time



Luminosities in PbA at $\sqrt{s}_{NN} = 72 \text{ GeV}$

With bent crystal

- ✓ Instantaneous luminosity: $L = \phi_{\text{beam}} \times N_{\text{target}} = \phi_{\text{beam}} \times (\rho \times l \times N_A) / A$
 l is a target thickness
- ✓ $\phi_{\text{beam}} = 2 \times 10^5 \text{ Pb s}^{-1}$
- ✓ Integrated luminosity - LHC year – 1 month running = 10^6 s


Target	$\rho \text{ (g.cm}^{-3}\text{)}$	A	L (mb ⁻¹ s ⁻¹)	$\int L \text{ (nb}^{-1}\text{yr}^{-1}\text{)}$
Liq H ₂ (1m)	0.07	1	800	800
Liq D ₂ (1m)	0.16	2	1000	1000
Be (1cm)	1.85	9	25	620
Cu (1cm)	8.96	64	17	17
W (1cm)	19.1	185	13	13
Pb (1cm)	11.35	207	7	7

- Nominal LHC luminosity for PbPb 0.5 nb^{-1}



Physics Highlights: AFTER@LHC

- *Physics Reports 522 (2013) 239; Few Body Syst. 53 (2012) 11-25.*
- *Many more ideas for a fixed target experiment at LHC submitted to a Special Issue in Advances in High*



Physics opportunities of a fixed-target experiment using LHC beams

S.J. Brodsky^a, F. Fleuret^b, C. Hadjidakis^c, J.P. Lansberg^{c,*}

^a SLAC National Accelerator Laboratory, Stanford University, Menlo Park, CA 94025, USA

^b Laboratoire Leprince Ringuet, Ecole polytechnique, CNRS/IN2P3, 91128 Palaiseau, France

^c IPNO, Université Paris-Sud, CNRS/IN2P3, 91406 Orsay, France

Contents

1. Introduction.....	6. Deconfinement in heavy-ion collisions.....
2. Key numbers and features.....	6.1. Quarkonium studies.....
3. Nucleon partonic structure.....	6.2. Jet quenching.....
3.1. Drell-Yan.....	6.3. Direct photon.....
3.2. Gluons in the proton at large x	6.4. Deconfinement and the target rest frame.....
3.2.1. Quarkonia.....	6.5. Nuclear-matter baseline.....
3.2.2. Jets.....	7. W and Z boson production in pp , pd and pA collisions.....
3.2.3. Direct/isolated photons.....	7.1. First measurements in pA
3.3. Gluons in the deuteron and in the neutron.....	7.2. W/Z production in pp and pd
3.4. Charm and bottom in the proton.....	8. Exclusive, semi-exclusive and backward reactions.....
3.4.1. Open-charm production.....	8.1. Ultra-peripheral collisions.....
3.4.2. $J/\psi + D$ meson production.....	8.2. Hard diffractive reactions.....
3.4.3. Heavy-quark plus photon production.....	8.3. Heavy-hadron (diffractive) production at $x_F \rightarrow -1$
4. Spin physics.....	8.4. Very backward physics.....
4.1. Transverse SSA and DY.....	8.5. Direct hadron production.....
4.2. Quarkonium and heavy-quark transverse SSA.....	9. Further potentialities of a high-energy fixed-target set-up.....
4.3. Transverse SSA and photon.....	9.1. D and B physics.....
4.4. Spin asymmetries with a final state polarization.....	9.2. Secondary beams.....
5. Nuclear matter.....	9.3. Forward studies in relation with cosmic shower.....
5.1. Quark nPDF: Drell-Yan in pA and Pbp	10. Conclusions.....
5.2. Gluon nPDF.....	Acknowledgments.....
5.2.1. Isolated photons and photon-jet correlations.....	References.....
5.2.2. Precision quarkonium and heavy-flavour studies.....	
5.3. Color filtering, energy loss, Sudakov suppression and hadron break-up in the nucleus.....	

Advances in High Energy Physics
 Hindawi
 Special Issue on
 Physics at a Fixed-Target Experiment Using the LHC
 Beams

CALL FOR PAPERS

http://after.in2p3.fr/after/index.php/Recent_published_ideas_in_favour_of_AFTER@LHC


- *Heavy-ion physics*
- *Exclusive reactions*
- *Spin physics studies*
- *Hadron structure*
- *Feasibility study and technical ideas*



Physics Highlights: AFTER@LHC

pp and pA @ $\sqrt{s_{NN}} = 115 \text{ GeV}$

- Understand dynamic of large-x gluon in nucleon
 - Quarkonia, Isolated photons, High- p_T jets ($> 20 \text{ GeV}/c$)
- ✓ Gluon distribution function in the proton: very large uncertainty at large x_B , also at large Q
- ✓ Unknown for the neutron





Physics Highlights: AFTER@LHC

pp and pA @ $\sqrt{s_{NN}} = 115 \text{ GeV}$

→ Understand dynamic of large-x gluon in nucleon

→ Quarkonia, Isolated photons, High- p_T jets ($> 20 \text{ GeV}/c$)

✓ Gluon distribution function in the proton: very large uncertainty at large x, also at large Q

✓ Unknown for the neutron

→ Heavy-quark distribution at large x




→ Open charm and beauty

✓ Pin down intrinsic charm

• Intrinsic heavy quarks are rigorous features of QCD

• Different charm pdfs (DGLAP or models with intrinsic charm) are in agreement with DIS data

*See also: arXiv:1504.06287,
arXiv:1410.0404, arXiv:0707.4658*



*CTEQ6.5C with
intrinsic charm*

Pumplin et al. Phys.Rev. D75 (2007)



Physics Highlights: AFTER@LHC

pp and pA @ $\sqrt{s_{NN}} = 115 \text{ GeV}$

→ Understand dynamic of large-x gluon in nucleon

→ Quarkonia, Isolated photons, High- p_T jets ($> 20 \text{ GeV}/c$)

✓ Gluon distribution function in the proton: very large uncertainty at large x, also at large Q

✓ Unknown for the neutron

→ Heavy-quark distribution at large x

→ Open charm and beauty

✓ Pin down intrinsic charm

• Intrinsic heavy quarks are rigorous features of QCD





• Different charm pdfs (DGLAP or models with intrinsic charm) are in agreement with DIS data

➤ With AFTER@LHC

• Good coverage in the target rapidity region

• High luminosity to reach large x_B

• Different targets: hydrogen, deuteron (neutron)



CTEQ6.5C with intrinsic charm

Pumplin et al. Phys.Rev. D75 (2007)



Physics Highlights: AFTER@LHC

pp and pA @ $\sqrt{s_{NN}} = 115 \text{ GeV}$


✓ Nucleon partonic structure

- Gluon pdf in the proton – large uncertainties at high x

- $g_p(x) = g_n(x)$?

 - Measure: quarkonia, isolated photons, high- p_T jets

 - Multiple probes to check factorization



✓ Heavy-quark distribution at large x in the proton

 - Measure: open heavy flavours


✓ Spin physics

- Gluon Sivers effect

- Linearly polarized gluons: $h_1^{\perp g}$, “Boers-Mulder” effect

- Single Spin Asymmetry in DY and HF studies

*See also: arXiv:1502.04021;
arXiv:1504.03791; arXiv:1504.04332,
arXiv:1203.5579; arXiv:1208.364*



✓ W and Z production near threshold ?

With AFTER@LHC: boost – better access to the low- p_T C-even quarkonia




Physics Highlights: AFTER@LHC

PbA @ $\sqrt{s_{NN}} = 72$ GeV, pA @ $\sqrt{s_{NN}} = 115$ GeV

→ Glue distribution in nucleus at large x

- Quarkonia
- Isolated photons
- High- p_T jets (> 20 GeV/c)

- ✓ Large uncertainty in nuclei at large x, unknown gluon EMC effect
- ✓ With AFTER@LHC:
 - Access to target $x_g = 0.3 - 1$ (>1 Fermi motion in nucleus)
 - With different targets:
 - probing A dependence of shadowing and nuclear matter effects





Physics Highlights: AFTER@LHC

PbA @ $\sqrt{s_{NN}} = 72 \text{ GeV}$

✓ Gluon distribution in nucleus at large x

- Complementary to EIC, LHeC
 - Quarkonia, isolated photons, high- p_T jets

✓ Quark-Gluon Plasma

• Experimental probes


- Quarkonia
- HF jets quenching
- Low mass lepton pairs
- Direct photons

- (Sequential ?) suppression of different quarkonium states – good resolution needed

- In PbA, different nuclei, A-dependent studies

- Precise estimation of Cold Nuclear Matter effects from pA

✓ Ultra-peripheral collisions





First simulations

Feasibility studies for quarkonium production at a fixed-target experiment using the LHC proton and lead beams (AFTER@LHC)

L. Massacrier,^{1,2} B. Trzeciak,³ F. Fleuret,⁴ C. Hadjidakis,² D. Kikola,⁵ J.P. Lansberg,² and H.-S. Shao⁶

¹*LAL, Université Paris-Sud, CNRS/IN2P3, F-91406, Orsay, France*

²*IPNO, Université Paris-Sud, CNRS/IN2P3, F-91406, Orsay, France*

³*FNSPE, Czech Technical U., Prague, Czech Republic*

⁴*Laboratoire Leprince Ringuet, École Polytechnique, CNRS/IN2P3, 91128 Palaiseau, France*

⁵*Faculty of Physics, Warsaw University of Technology, ul. Koszykowa 75, 00-662 Warsaw, Poland*

⁶*PH Department, TH Unit, CERN, CH-1211, Geneva 23, Switzerland*

(Dated: June 17, 2015)

Used in the fixed-target mode, the multi-TeV LHC proton and lead beams allow for studies of heavy-flavour hadroproduction with unprecedented precision at backward rapidities –far negative Feynman- x – using conventional detection techniques. At the nominal LHC energies, quarkonia can be studied in detail in $p + p$, $p + d$ and $p + A$ collisions at $\sqrt{s_{NN}} \approx 115$ GeV as well as in $Pb + p$ and $Pb + A$ collisions at $\sqrt{s_{NN}} \approx 72$ GeV with luminosities roughly equivalent to that of the collider mode, *i.e.* up to $20 \text{ fb}^{-1}\text{yr}^{-1}$ in $p + p$ and $p + d$ collisions, up to $0.6 \text{ fb}^{-1}\text{yr}^{-1}$ in $p + A$ collisions and up to $10 \text{ nb}^{-1}\text{yr}^{-1}$ in $Pb + A$ collisions. In this paper, we assess the feasibility of such studies by performing fast simulations using the performance of a LHCb-like detector.

arXiv: 1504.5145





Advances in High Energy Physics

Special Issue on
**Physics at a Fixed-Target Experiment Using the LHC
Beams**

CALL FOR PAPERS



Charge particle multiplicities in a fixed target mode



- ✓ Charge particle multiplicities, for all possible fixed target modes, p+Pb, Pb+H, Pb+Pb, are smaller than the ones reached in the collider modes. A detector with the LHCb capabilities will be able to run in such conditions (LHCb was used in p+Pb and Pb+p at 5 TeV).



Expected quarkonium yield

pp and pA @ $\sqrt{s} = 115 \text{ GeV}$

Target	$\int \mathcal{L} \text{ (fb}^{-1}\cdot\text{yr}^{-1}\text{)}$	$N(\text{J}/\Psi) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Upsilon}$
1 m Liq. H ₂	20	4.0 10 ⁸	8.0 10 ⁵
1 m Liq. D ₂	24	9.6 10 ⁸	1.9 10 ⁶
LHC pp 14 Tev (low pT)	0.05 (ALICE) 2 LHCb	3.6 10 ⁷ 1.4 10 ⁹	1.8 10 ⁵ 7.2 10 ⁶
RHIC pp 200GeV	1.2 10 ⁻²	4.8 10 ⁵	1.2 10 ³

pp

1 m H₂ target

- ✓ 1000 times more statistics than at RHIC (@200 GeV)
- ✓ Comparable statistics to LHC

pA

Target	A	$\int \mathcal{L} \text{ (fb}^{-1}\cdot\text{yr}^{-1}\text{)}$	$N(\text{J}/\Psi) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= A\mathcal{L}B\sigma_{\Upsilon}$
1cm Be	9	0.62	1.1 10 ⁸	2.2 10 ⁵
1cm Cu	64	0.42	5.3 10 ⁸	1.1 10 ⁶
1cm W	185	0.31	1.1 10 ⁹	2.3 10 ⁶
1cm Pb	207	0.16	6.7 10 ⁸	1.3 10 ⁶
LHC pPb 8.8 TeV	207	10 ⁻⁴	1.0 10 ⁷	7.5 10 ⁴
RHIC dAu 200GeV	198	1.5 10 ⁻⁴	2.4 10 ⁶	5.9 10 ³
RHIC dAu 62GeV	198	3.8 10 ⁻⁶	1.2 10 ⁴	18

1 cm Pb target

- ✓ 100 times more statistics than at RHIC (dAu@200 GeV)
- ✓ Comparable statistics to LHC

Detailed study of quarkonium production and nuclear effects

See also: *Advances in High Energy Physics*, Article ID 726393, in press.
arXiv:1504.0653



Expected quarkonium yield

$$\text{PbA} @ \sqrt{s}_{\text{NN}} = 72 \text{ GeV}$$

PbA

Target	A.B	$\int \mathcal{L} \text{ (nb}^{-1}\cdot\text{yr}^{-1}\text{)}$	$N(\text{J}/\Psi) \text{ yr}^{-1}$ $= \text{AB} \mathcal{L} \mathcal{B} \sigma_{\Psi}$	$N(\Upsilon) \text{ yr}^{-1}$ $= \text{AB} \mathcal{L} \mathcal{B} \sigma_{\Upsilon}$
1 m Liq. H ₂	207.1	800	3.4 10⁶	6.9 10³
1cm Be	207.9	25	9.1 10⁵	1.9 10³
1cm Cu	207.64	17	4.3 10⁶	0.9 10³
1cm W	207.185	13	9.7 10⁶	1.9 10⁴
1cm Pb	207.207	7	5.7 10⁶	1.1 10⁴
LHC PbPb 5.5 TeV	207.207	0.5	7.3 10⁶	3.6 10⁴
RHIC AuAu 200GeV	198.198	2.8	4.4 10⁶	1.1 10⁴
RHIC AuAu 62GeV	198.198	0.13	4.0 10⁴	61

1 cm Pb target

- ✓ Similar statistics than at RHIC @200 GeV
- ✓ 2 order of magnitude larger than at RHIC @62 GeV

Detailed study of
quarkonium states



First simulations of quarkonia, pp at $\sqrt{s} = 115$ GeV


- PYTHIA 8.185, fast simulations with LHCb-like reconstruction parameters

- ✓ Requirements:

- ➔ Momentum resolution: $\Delta p/p = 0.5\%$
- ➔ μ identification efficiency: 98%

- ✓ Single μ cuts:




- ➔ $2 < \eta_\mu < 5$
- ➔ $p_T^\mu > 0.7$ GeV/c
- ➔ μ misidentification (with π or K)



- ✓ Input for quarkonium signals: HELAC-Onia

- ✓ Estimation of different dimuon background sources:

- ➔ Uncorrelated background – min bias PYTHIA 8
- ➔ Drell-Yan – HELAC-Onia
- ➔ cc, bb – HELAC-Onia



Efficiency of background μ pairs




ψ and Υ signal simulations with full background


$$J/\psi / \psi(2S) \rightarrow \mu^+ \mu^-$$

$$\Upsilon(nS) \rightarrow \mu^+ \mu^-$$

$\int L = 10 \text{ fb}^{-1}$, **0.5 year of data taking with 1m H₂ target** (in the crystal case)



→ Dominant source of background is uncorrelated background





→ Dominant source of background is DY
→ Clear separation of different states

J/ψ signal simulation with full background

$$J/\psi \rightarrow \mu^+ \mu^-$$



$\int L = 10 \text{ fb}^{-1}$, **0.5 year of data taking with 1m H₂ target** (in the crystal case)



- p_T and rapidity Distributions for the J/ψ and different backgrounds differ.
- In more backward or forward rapidity regions, the signal to background ratio increases

Quarkonium acceptance and p_T reach

$\int L = 10 \text{ fb}^{-1}$, **0.5 year of data taking with 1m H₂ target** (in the crystal case)




- J/ψ and ψ(2S) signals can be studied up to $\sim 15 \text{ GeV}/c$, $\Upsilon(nS)$ up to $\sim 10 \text{ GeV}/c$
- All quarkonium states can be measured down to $0 \text{ GeV}/c$
- Similar p_T reach expected for pA



- Study is limited to the rapidity range of $2 < y < 5$ ($2 < \eta_\mu < 5$)
- J/ψ and ψ(2S) signals can be studied in the whole range, lowest y for $\Upsilon(nS)$ is $\sim 2.5-3$

Impact of nPDF effects on quarkonium R_{pPb}


→ Combination of measurements of $\Upsilon(nS)$, J/ψ and $\psi(2S)$ for $-3 < y_{CMS} < 0$ (as LHCb detector would do) will allow to pin down the existence of a possible gluon EMC and antishadowing effect



Simulations done using JIN with EPS09



Very good statistical precision !



See also: *Advances in High Energy Physics*, Article ID 492302 and 783134, in press; [arXiv:1507.05413](https://arxiv.org/abs/1507.05413); [arXiv:1504.07428](https://arxiv.org/abs/1504.07428)

pp: $\int L = 10 \text{ fb}^{-1}$, pPb: $\int L = 100 \text{ pb}^{-1}$



Summary

after.in2p3.fr

- Many physics opportunities with a fixed target experiment using LHC p and Pb beams
- Novel testing ground for QCD in the high-x frontier with AFTER@LHC
- Extensive spin program with a polarized target
- Using dense targets high luminosities can be achieved
- Target versatility: hydrogen, deuteron, nucleus – nuclear effects and QGP
- First fast simulations performed
- ➔ Simulations in pA, AA and of different quarkonium states in progress

Thank you !

This work was supported by the European social fund within the framework of realizing the project „Support of inter-sectoral mobility and quality enhancement of research teams at Czech Technical University in Prague“, CZ.1.07/2.3.00/30.0034.



BACKUP




Outlook

- Special Issue in Advances in High Energy Physics
- Expression of interest expected in 2015/2016
- Development of the fast simulation framework

after.in2p3.fr

Advances in High Energy Physics



Special Issue on
Physics at a Fixed-Target Experiment Using the LHC Beams

CALL FOR PAPERS

Fixed-target experiments (FTE) have brought essential contributions to particle and nuclear physics. They have led to particle discoveries (Ω , J/ψ , ...) and evidence for the novel dynamics of quarks and gluons in heavy-ion collisions. In accessing high x_F and in offering options for (un-) polarised proton and nuclear targets, they have also led to the observation of surprising QCD phenomena. They offer specific advantages compared to collider experiments: access to high x_F , high luminosities, target versatility, and polarisation.

The LHC 7 TeV protons on targets release a c.m.s. energy close to 115 GeV (72 GeV with Pb), in a range never explored so far, significantly higher than that at SPS and not far from RHIC. The production of quarkonia, DY, heavy flavours, jets, and γ in pA collisions can be studied with statistics previously unheard of and in the backward region, $x_F < 0$, which is uncharted. High precision QCD measurements can also obviously be carried out in pp and pA collisions with H_2 and D_2 targets. With the 50 TeV protons of the future circular collider (FCC), the c.m.s. energy could reach 300 GeV for original studies of W and Z boson, and perhaps H^0 , production in pp and pA collisions.

With the LHC Pb beam, one can study the quark-gluon plasma (QGP) from the viewpoint of the nucleus rest frame after its formation. Thanks to modern technologies, studies of, for instance, direct γ and quarkonium P -waves production in heavy-ion collisions can be envisioned.

Polarising the target allows one to study single-sptn correlations including the Sivers effect, hence, the correlation between the parton k_T and the nucleon sptn.

We intend to publish a special issue on the physics at such a FTE using the LHC or FCC beams. The editors welcome original research articles and review articles from both theorists and experimentalists.

Potential topics include, but are not limited to:

- ▶ Heavy-quark and gluon content at large x
- ▶ TMDs and single-sptn asymmetries
- ▶ Heavy-flavour studies in pA and AA collisions at FTEs
- ▶ W, Z, and H^0 production near threshold
- ▶ Target polarisation
- ▶ Secondary beams
- ▶ Simulation tools for high-energy physics
- ▶ Beam collimation and extraction with bent crystals
- ▶ Machine feasibility and radiological aspects
- ▶ Connection between UHECR studies and FTEs

Lead Guest Editor
Jean-Philippe Lansberg, IPN Orsay, Orsay, France
jlansberg@in2p3.fr

Guest Editors
Gianluca Cavoto, Istituto Nazionale Di Fisica Nucleare, Roma, Italy
gianluca.cavoto@roma1.infn.it

Cynthia Hadjiidakis, IPN Orsay, Orsay, France
cynthia@ipno.in2p3.fr

Jibo He, CERN, Geneva, Switzerland
jibo.he@cern.ch

Cédric Lorcé, Université de Liège, Liège, Belgium
c.lorce@ulg.ac.be

Barbara Trzeciak, Czech Technical University, Prague, Czech Republic
trzeciak@fffi.cvut.cz

Manuscript Due
Friday, 20 March 2015


First Round of Reviews
Friday, 12 June 2015

Publication Date
Friday, 7 August 2015

Beam extraction using bent crystal




✓ Possible fixed-target mode




W. Scandale et al., JINST 6 T10002 (2011)

UA9 experiment @ SPS, 15/10/2014



Direct view of the channeled beam



S. Montesano, W. Scandale, Joint LUA9-AFTER meeting, Nov. 2013



Beam extraction using bent crystal

✓ Beam collimation @LHC: amorphous collimator, inefficiency of 0.2% (3.5 TeV p beam)

• Expected bent crystal inefficiency: 0.02%

UA9: test @SPS on the crystal with proton and ion beams

LUA9 (beam bending experiment using crystal): approved by LHCC

2 bent crystals installed in IR7 during LS1

2015/2016 first tests with beams

➤ Proton beam extraction:

• Single or multi-pass extraction efficiency of 50%

• LHC beam loss $\sim 10^9 \text{ p}^+ \text{ s}^{-1}$ - extracted beam : $5 \times 10^8 \text{ p}^+ \text{ s}^{-1}$

• Extremely small emittance: beam size (in the extraction direction) 950m after the extraction: 0.3mm

➤ Ion beam extraction

• Successfully tested at the SPS, should also work at the LHC (P. Ballin et al, NIMB 267 (2009) 2952)

➔ Deflecting the beam halo at 7σ distance to the beam

➔ No loss in the LHC beam



Physics Highlights: AFTER @ LHC

pp and pA @ $\sqrt{s_{NN}} = 115 \text{ GeV}$

(Gluon) Sivers effects with a transversely polarized target

Gluon Sivers effect: correlation between the gluon transverse momentum k_T and the proton spin

- The target rapidity region ($x_F < 0$) corresponds to high x^\uparrow ($x_F \rightarrow -1$) where the k_T - spin correlation is the largest
- Transverse single spin asymmetries studied using **gluon sensitives probes**:
 - quarkonia (J/ψ , Υ , χ_c)
 - B & D mesons production
 - γ , γ -jet, γ - γ also J/ψ - γ

L. Massacrier – SPIN 2014
Conference



Physics Highlights: AFTER @ LHC

pp and pA @ $\sqrt{s}_{NN} = 115 \text{ GeV}$

TMDs STUDIES WITH AFTER@LHC (WITH A POLARIZED TARGET)


(Quark) Sivers effects with a transversely polarized target

□ Can be probed with the Drell-Yan

Experiment	particles	energy (GeV)	\sqrt{s} (GeV)	x_p^\dagger	\mathcal{L} ($\text{nb}^{-1}\text{s}^{-1}$)
AFTER	$p + p^\dagger$	7000	115	$0.01 \div 0.9$	1
COMPASS	$\pi^\pm + p^\dagger$	160	17.4	$0.2 \div 0.3$	2
COMPASS (low mass)	$\pi^\pm + p^\dagger$	160	17.4	~ 0.05	2
RHIC	$p^\dagger + p$	collider	500	$0.05 \div 0.1$	0.2
J-PARC	$p^\dagger + p$	50	10	$0.5 \div 0.9$	1000
PANDA (low mass)	$\bar{p} + p^\dagger$	15	5.5	$0.2 \div 0.4$	0.2
PAX	$p^\dagger + \bar{p}$	collider	14	$0.1 \div 0.9$	0.002
NICA	$p^\dagger + p$	collider	20	$0.1 \div 0.8$	0.001
RHIC	$p^\dagger + p$	250	22	$0.2 \div 0.5$	2
Int.Target 1					
RHIC Int.Target 2	$p^\dagger + p$	250	22	$0.2 \div 0.5$	60
P1027	$p^\dagger + p$	120	15	$0.35 \div 0.85$	400-1000
P1039	$p + p^\dagger$	120	15	$0.1 \div 0.3$	400-1000

Relevant parameters for the future proposed polarized DY experiments

S. J. Brodsky et al., Phys. Rep. 522 (2013) 239
 V. Barone et al., Prog. Part. Nucl. Phys. 65 (2010) 267



Prediction for AFTER

M. Anselmo, ECT*, Feb. 2013
 (Courtesy U. d'Alessio)

Asymmetry up to 10% predicted in DY for the target rapidity region ($x_F < 0$)

L. Massacrier – SPIN 2014
 Conference



Physics Highlights: AFTER@LHC


pp and pA @ $\sqrt{s_{NN}} = 115 \text{ GeV}$

✓ Linearly polarized gluons: $h_1^{\perp g}$

→ “Boers-Mulder” effect: correlation between the parton k_T and its spin (in unpolarized nucleon)

→ Scalar and pseudo-scalar quarkonia – χ_{c0} ,

χ_{b0} , η_c , η_b



✓ Low- p_T C-even quarkonium production is a good probe of gluon Transverse Momentum Dependent (TMD) pdfs


✓ Low- p_T scalar and pseudo-scalar quarkonia are affected differently by the linearly polarized gluons in unpolarized nucleons

✓ With AFTER@LHC

• Boost – better access to the low- p_T C-even quarkonia




• η_c (LHCb 1409.3612), (η_b), back-to-back $J/\psi + \gamma$, $J/\psi + J/\psi$

Boer, Pisano, PRD 86 (2012) 094007



ψ signal simulation with full background

$$J/\psi / \psi(2S) \rightarrow \mu^+ \mu^-$$



- In more backward or forward rapidity regions, the signal to background ratio increases